



اللجنة الأكاديمية للهندسة المدنية

ملخص

لاب كيمياء

نمر عودة

Contact us:

f Civilittee HU | لجنة المدني

▶ Civilittee Hashemite

www.civilittee-hu.com



Exp 1 Safety and Equipment. There is a quiz for the exp.

المطلوب:- فقط الأدوات صفحة 4-7 + فقط قواعد السلامة من 1-13 من
 الترتيب حسب الدقة + فقط أدوات السلامة ص 8+9
 فقط volumetric pipette & volumetric flask > pipette & Burette > graduated cylinder > Beaker & Er. Flask

Exp 2 Empirical Formula of a compound. There is a quiz for the exp.

* هذه التجربة تحتوي جانب نظري وجانب حسابي
 * النظري يتعلل بالعلاقات والحسابي يتعلل بالتحويلات بناء على الجدول الدوري
 كما في المادة الأصلية (أساسيات الكيمياء)

Calculation Part:-

Ex) How many grams of Magnesium combine with 1.5g of chloride ions in $MgCl_2$?

Sol →

المطلوب: كم في المادة النظرية

$$1.5g Cl \times \frac{1 \text{ mol Cl}}{35.4g Cl} \times \frac{1 \text{ mol MgCl}}{2 \text{ mol Cl}} \times \frac{24.31g Mg}{1 \text{ mol Mg}} = 0.514g Mg$$

Ex) IF 11.8 g of iron reacts with 5.06 g of Oxygen. Determine the empirical formula of the resulting oxide?

Sol →

الخطوات

Fe : O

11.8g : 5.06g

① تحويل لغرامات

$$\frac{11.8}{55.8} = 0.211 \text{ mol} : \frac{5.06g}{16} = 0.316 \text{ mol}$$

② تحويل لمولات


$$\frac{0.211}{0.211} = 1 : \frac{0.316}{0.211} = 1.5$$

③ القسمة على أصغر عدد

$$1 \times 2 = 2$$

$$1.5 \times 2 = 3$$

④ أكبر الرقم



Ex) Nicotine is a compound containing C, H and N. A 2.5 g sample of the compound is burned and produces 6.78 g of CO_2 , 1.94 g of H_2O , and 0.43 g of N_2 . What is the empirical formula of nicotine?

Sol →

$$\begin{array}{lcl}
 \text{C} & \text{H} & \text{N} \\
 6.78 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44 \text{ g CO}_2} \times \frac{1 \text{ mol C}}{1 \text{ mol CO}_2} & 1.94 \text{ g H}_2\text{O} \times \frac{1 \text{ mol H}_2\text{O}}{18 \text{ g H}_2\text{O}} \times \frac{2 \text{ mol H}}{1 \text{ mol H}_2\text{O}} & 0.43 \text{ g N}_2 \times \frac{1 \text{ mol N}_2}{28 \text{ g N}_2} \times \frac{2 \text{ mol N}}{1 \text{ mol N}_2} \\
 = 0.154 \text{ mol} & = 0.216 \text{ mol} & = 0.031 \text{ mol}
 \end{array}$$

① داجار مولان كل عنصر

$$0.154 / 0.031 = 5 \quad ; \quad 0.216 / 0.031 = 7 \quad ; \quad 0.031 / 0.031 = 1$$

② النسبة على الصغرى



Ex) When 0.288 g of P is burned, 0.66 g of white Phosphorus oxide is obtained, determine the empirical formula of this oxide?

Sol → we have 0.288 g P & 0.66 g of Oxide

$$\begin{aligned}
 \text{Mass of O} &= \text{Mass of Oxide} - \text{Mass of P} = 0.66 - 0.288 \\
 &= 0.372 \text{ g}
 \end{aligned}$$



① $0.288 \text{ g} : 0.372 \text{ g}$

② $\frac{0.288}{31} = 0.0093 \text{ mol} ; \frac{0.372}{16} = 0.0233 \text{ mole}$

③ $\frac{0.0093}{0.0093} = 1 ; \frac{0.0233}{0.0093} = 2.5$

④ $1 \times 2 = 2 ; 2.5 \times 2 = 5$



Ex) A 2.00-g Sample of a bromide oxide is converted to 2.936 g of AgBr, Calculate the empirical formula of the oxide, (M.wt for AgBr = 187.78 g/mol, Br = 80 g/mol and O = 16 g/mol)

Sol → نفس فكرة السؤال السابق لكن يجب أولاً إيجاد كتلة Br

$$2.936 \text{ g AgBr} \times \frac{1 \text{ mol AgBr}}{187.78 \text{ g AgBr}} \times \frac{1 \text{ mol Br}}{1 \text{ mol AgBr}} \times \frac{80 \text{ g Br}}{1 \text{ mol Br}} = 1.25 \text{ g Br}$$

$$\text{mass O} = \text{mass Oxide} - \text{mass Br} = 2 \text{ g} - 1.25 \text{ g} = 0.75 \text{ g O}$$

Br : O

$$\frac{1.25}{80} = 0.0156 \text{ mol} ; \frac{0.75}{16} = 0.0468$$

$$\frac{0.0156}{0.0156} = 1 ; \frac{0.0468}{0.0156} = 3$$

BrO₃

* أسئلة الحسابات في هذه التجربة لا تنقد الأفكار السابقة

The Other Part :-

① أسئلة تتعلق بالتجربة :-

why?

- [1] Heating before starting? To remove the moisture الرطوبة
- [2] Don't weight the crucible when it's hot? It gives wrong accurate
- [3] Don't cover the crucible widely? It burns Mg brightly
- [4] Adding a few water drops? To decompose Mg₃N₂

Postlab questions

③

What is the effect of Mg:O mole ratio on:

- ① Mg_3N_2 Not decomposed completely $\rightarrow$ increased
 $\nwarrow$ OR The formation of side product.
 - ② Mg_3N_2 decomposed completely $\rightarrow$ no effect.
 - ③ Carbon deposited on the crucible surface $\rightarrow$ decrease
 - ④ Carbon not deposited on the crucible surface $\rightarrow$ no effect
 - ⑤ Magnesium oxide ash is not dried completely $\rightarrow$ decrease
 - ⑥ Magnesium oxide ash is dried completely $\rightarrow$ no effect
 - ⑦ Rapid Oxidation of Magnesium $\rightarrow$ increase
 - ⑧ Air is not sufficient to ~~react~~^{react} with all the Mg $\rightarrow$ increase
 - ⑨ Air is sufficient to react with all the Mg $\rightarrow$ no effect
 - ⑩ Nonvolatile and unreactive impurities in the crucible during oxidation $\rightarrow$ decrease
 - ⑪ Nonvolatile and unreactive impurities in the crucible from the beginning $\rightarrow$ no effect
 - ⑫ If the balance reads $\pm 0.02g$ for any reading $\rightarrow$ no effect
- * بالنسبة ل 10 + 11 باختصار $\rightarrow$ ① و ② مواد متطايرة قبل التأكسد لا يؤثر وأثناء التأكسد يؤثر.
 ③ وجود مواد متطايرة قبل أو أثناء التأكسد لا يؤثر.
 (dec)

* لو طلب O:Mg $\rightarrow$ نكتب العلاقات أعلاه

Eq $\rightarrow$ بقا

- ① The formation of Mg_3N_2 (side reaction) $\rightarrow 3Mg + N_2 \rightarrow Mg_3N_2$
- ② The reaction between magnesium Nitride and water $\rightarrow Mg_3N_2 + 6H_2O \rightarrow 3Mg(OH)_2 + 2NH_3$
- ③ Heating Mg to ash $\rightarrow Mg(OH)_2 \xrightarrow{\Delta} MgO + H_2O$

Exp 3 → Limiting Reactant.
* No quiz for the exp.

Page 5

* الهدف من هذه التجربة هو معرفة كيفية تحديد ال L.R

⊗ Calculation Part →


$$\text{Percent yield} = \frac{\text{Actual yield} \xrightarrow{\text{given}}}{\text{Theoretical yield} \xrightarrow{\text{calculated}}} \times 100\%$$

= How to determine the theoretical yield?

- ① we find the L.R by →
 - [1] convert each element to moles
 - [2] Divide each element on it's coefficient
 - [3] The smallest number is the L.R

② we use only the L.R in our normal calculation.


* Before we start -



* التجربة سوف تقوم على إجراء هذه المعادلة هي حفظ

Ex) A 25 g sample of $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ react with excess $\text{BaCl}_2 \cdot \text{H}_2\text{O}$. If the mass of $\text{Ba}_3(\text{PO}_4)_2$ obtained is 17.56g. Calculate the % yield of $\text{Ba}_3(\text{PO}_4)_2$?

Sol →



[1] Determine the L.R → given Ba^{+2} excess so PO_4^{-3} is L.R

[2] Use grams of PO_4^{-3} in our normal calculations:-

$$\text{Theoretical yield} = 25 \text{ g } \cancel{\text{PO}_4^{-3}} \times \frac{1 \text{ mol } \cancel{\text{PO}_4^{-3}}}{380.2 \text{ g } \cancel{\text{PO}_4^{-3}}} \times \frac{1 \text{ mol } \text{Ba}_3(\text{PO}_4)_2}{2 \text{ mol } \cancel{\text{PO}_4^{-3}}} \times \frac{602.2 \text{ g } \text{Ba}_3(\text{PO}_4)_2}{1 \text{ mol } \text{Ba}_3(\text{PO}_4)_2}$$

$\xleftarrow{\text{from the eq}}$

$$\text{Actual yield} = 17.56 \rightarrow \% \text{ yield} = \frac{17.56}{19.8} = 88.68\%$$

Ex) A mixture containing 40 g of $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ and 30 g of $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ is dissolved in water. A precipitate of $\text{Ba}_3(\text{PO}_4)_2$ weighing 22.65 g is produced. Calculate the % yield of $\text{Ba}_3(\text{PO}_4)_2$.

↳ actual yield.

Sol →

[1] Determine the L.R

$$\frac{40 \text{ g } \text{PO}_4^{3-}}{380.2} = 0.105 \text{ mole} \rightarrow \frac{0.105 \text{ mole}}{\text{coefficient } \leftarrow 2} : 0.0525$$

$$\frac{30 \text{ g } \text{Ba}^{+2}}{244.2} = 0.123 \text{ mole} \rightarrow \frac{0.123 \text{ mole}}{3} = 0.041 \rightarrow \text{The Smallest}$$

So Ba^{+2} is L.R

[2]

$$0.123 \text{ mol } \text{Ba}^{+2} \times \frac{1 \text{ mol } \text{Ba}_3(\text{PO}_4)_2}{3 \text{ mol } \text{Ba}^{+2}} \times \frac{602.7 \text{ g } \text{Ba}_3(\text{PO}_4)_2}{1 \text{ mol } \text{Ba}_3(\text{PO}_4)_2}$$


Theoretical

~~Actual~~ yield = 24.69 g

% yield = $\frac{22.65}{24.69} \times 100\% = 91.7\%$

Ex) 10 g of a unknown mixture containing $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ and $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ is dissolved in distilled water. The mass of $\text{Ba}_3(\text{PO}_4)_2$ precipitated is 3.5 g. calculate the % of each salt present in the mixture. if the BaCl_2 is the limiting reactant?

[1] Ba^{+2} is L.R



we have given 3.5 g $\text{Ba}_3(\text{PO}_4)_2$ so

$$3.5 \text{ g } \text{Ba}_3(\text{PO}_4)_2 \times \frac{1 \text{ mol } \text{Ba}_3(\text{PO}_4)_2}{602.7 \text{ g } \text{Ba}_3(\text{PO}_4)_2} \times \frac{3 \text{ mol } \text{Ba}^{+2}}{1 \text{ mole } \text{Ba}_3(\text{PO}_4)_2} \times \frac{244.2 \text{ g } \text{Ba}^{+2}}{1 \text{ mol } \text{Ba}^{+2}}$$

Mass $\text{Ba}^{+2} = 4.25 \text{ g}$

% $\text{Ba}^{+2} = \frac{4.25 \text{ g}}{10 \text{ g}} \times 100\% = 42.5\%$

% $\text{PO}_4^{3-} = (100 - 42.5)\% = 57.5\%$


* Procedures Part →

why?

- 1] Don't boil the solution → To minimize the lost of the mass
- 2] Using distilled water → To provide the reactions of the ^{unknowns} ~~unknowns~~

* What is the supernatant liquid?

It is the clear liquid above a precipitate.

* يعني في طبقة السائل
 طبقة السائل
 lig. 
 Prec.

* Determination of the L.R →

1] Test for excess PO_4^{3-} OR Limiting Ba^{+2} →

Add 2 drops of 0.5M BaCl_2 to the solution. If a precipitate is formed then PO_4^{3-} is the excess and Ba^{+2} is the L.R. .
 If a precipitate is not formed then the PO_4^{3-} is the ^{L.R} ~~excess~~ and Ba^{+2} is the excess.

2] Test for excess Ba^{+2} OR Limiting PO_4^{3-} →

Add 2 drops of 0.5M Na_3PO_4 to the solution. If a ppt. is formed then Ba^{+2} is the excess and PO_4^{3-} is the L.R.
 If a ppt. is not formed then Ba^{+2} is the L.R and PO_4^{3-} is the excess.

Post Lab Part

≡ what is the effect of heating the solution on the particle size of $\text{Ba}_3(\text{PO}_4)_2$ ppt.? It makes a conglutination

≡ what is the effect on the actual yield of the $\text{Ba}_3(\text{PO}_4)_2$ if:-




- 1] Using a coarse paper → decrease
- 2] Insufficient washing of the ppt. → increase.
- 3] Using Acidic to wash the solution → decrease.
- 4] The ppt. wasn't dried completely → increase.

Expt 4




Tests For Cations and Anions

There is no quiz for the exp.

Cations →

[1] Ca^{+2} , Calcium test[2] Fe^{+3} , Ferric test[3] NH_4^+ ammonium test

Anions →

[1] SO_4^{-2} (sulfate test)[2] HCO_3^- - test[3] Test of Cl^- , Br^- 

* المطلوب من هذه التجربة هو فقط المكتوب أدناه .
 * المعادلات ليست مهمة لكن المطلوب أسماء المركبات وكيفية الكشف .
 والكواشف ولون الناتج والوسط الموضوع فيه فقط .

Exps Identification of A compound:

Page 9

Physical Properties.

There is no quiz for the exp.

* PreLab →

[1] Define

A. Solubility → ^{الذائبة} The maximum mass of solute that can be dissolved in a fixed mass of solvent at a given temperature.

B. Boiling Point → [1] The temperature at which the liquid starts to boil

[2] The temperature at which bubble form spontaneously and continue to form until the entire volume of the liquid has been converted to a gas.

[3] The temperature where the vapor pressure is equal to the atmospheric pressure

[2] A student's liquid unknown boils at approximately 69°C , is insoluble in water but soluble in cyclohexane. Its density is 0.69 g/ml , which chemical in table (page 41) is the unknown?

* هذا السؤال إذا أجاب يبي مع جدول، فاعليش، الاتقارن بين المعطيات والجدول وقتاً.

The unknown is → n-hexane.

[3] What physical property, measurable in this experiment, distinguishes cyclohexane from cyclohexene?

Boiling point.

[4] Using apparatus described in this experiment, when should the boiling point of a liquid be recorded?

When the bubbles cease to escape and before the liquid re-enters the capillary tube.

* The exp.

II Solubility → الذائبة

Solution = Solute + Solvent
محلول مذاب مذيب

* Likes dissolves in like

1 Polar - Polar $\rightarrow$ Soluble (miscible) يذوب

[2] non polar - non polar $\rightarrow$ Soluble (miscible) يزوب

[3] Polar - Salt $\rightarrow$ Soluble (miscible) ينوب

□ nonpolar - Salt → insoluble (immiscible) لا يذوب

5) Polar - nonpolar $\rightarrow$ insoluble (immiscible) لا يذوب

Levels of solubility $\rightarrow$

[1] complete dissolving [2] partial dissolving [3] insoluble.
 ذوبان کامل ذوبان جزئی غیر ذائب (عقیم).

[2] Partial dissolving [3] insoluble.
ذوبان جزئي غير ذائب (عديم).

خير ذائب (عديم).

② Density = $\frac{\text{mass}}{\text{Volume}}$, $\rho = \frac{m}{V}$ 1 L = 1000 mL

[3] Boiling point, B.P

it called normal B.P if the vapor pressure = $P_{atm} = 1$

B.P solution > B.P ~~Solute~~ Solvent why? $\rightarrow$ Because it has stronger intermolecular forces.

* we are dealing with 3 solvents in this experiment.

① Water (H_2O) → Polar

② Cyclohexane $\rightarrow$ $C_6H_{12} \rightarrow$ non-polar

③ Ethanol (C_2H_5OH) $\rightarrow$ Polar

* Post Lab →

II How does atmospheric pressure affect the boiling point of a liquid?

directly $\rightarrow$ $\vec{b}_1, \vec{b}_2$

* Also $\rightarrow$

How does Intermolecular forces affect the boiling point of a liquid?
directly

2] If several drops of liquid unknown ^{بالصق} cling to the pipette wall after delivery will the density of the unknown be reported too high or too low?

too low because mass will decrease then due to the relation $\rho = \frac{m}{V}$, the density will decrease.

3] A. If the ~~re~~ Boiling pt. is recorded when bubbles are rapidly escaping the capillary tube, will it be recorded too high or too low? Explain.

* إذا قرأنا بعد ما طلع الماء من الأنبوب ← القراءة < الأصلية

temp > true B.P, because $V.P > P_{atm}$

B. If the Boiling pt. is recorded after the liquids enters the capillary tube (after the heat is removed), will it be recorded too high or too low? Explain.

* إذا قرأنا بعد ما دخل الماء إلى الأنبوب ← القراءة > الأصلية

temp. < true B.P, because $V.P < P_{atm}$.

C. If the Boiling pt. is recorded when the liquids cease to escape and before the liquids re-enters the capillary tube?

* إذا قرأنا أثناء الخروج وقبل الدخول إلى الأنبوب ← القراءة = الأصلية

temp = true B.P, because $V.P = P_{atm}$.

* إذا أجبنا أي سؤال ببيكليت أي المواد الآتية تصلح للتجربة إختار، التي درجة غليانها أقل من 100 (أقل من درجة غليان الماء).

* إذا أجبنا أي سؤال ببيكليت أي المواد الآتية لا تصلح للتجربة إختار، التي درجة غليانها أكبر من 100 (أكبر من درجة غليان الماء).

Can you predict when the $V.P = P_{atm}$, theoretically? No

Exp 1

Q₁) The correct statement concerning handling of chemicals in Lab is:-

1. Direct contact with chemicals is allowed.
2. No need to read the label on the reagent bottles.
3. Smelling and tasting chemicals are not allowed.
4. Toxic chemicals can be used outside the fume hood.
5. All of the above.

Ans: 3

Q₂) Which of the following is not a safety tool?

1. First aid Equipment.
2. Fire extinguisher
3. Graduated cylinder
4. Fume hood
5. Fire blanket.

Ans: 3

Q₃) Which of the following is not a safety equipment?

1. Beaker 2. Goggle 3. First aid equipments 4. Fire blanket

Ans: 1

Q₄) Write down T or F:-

T Do not point your test tube at your face when heating anything to watch what is happening exactly.

F Open sandals, short skirts and shorts are allowed in the lab.

Q5) When a metal (M) with atomic mass 56 g/mol was oxidized to a metal oxide that contains 36.4% by mass O (Atomic weight O = 16 g/mol), the empirical formula of the metal oxide is:-

1. MO_2
2. MO_3
3. M_2O_3
4. M_3O_4
5. M_2O_5

Sol → 36.4 % O → 36.4 g O } تحويل الغرامات
 %M = (100 - 36.4)% = 63.6% → 63.3 g M

$$\begin{array}{r} M: \quad O \\ \hline 63.3 \text{ g} \quad 36.4 \text{ g} \\ 56 \quad 16 \\ \hline 1.13 \text{ mol} \quad 2.275 \text{ mol} \\ 1.13 \quad 1.13 \end{array}$$

تحويل المولات

النسبة إلى الأصغر

1 : 2

So MO_2

Ans: 1

Q6) In the empirical formula experiment which statement below is incorrect:-

1. (Mg to O) mole ratio will not affect, if the balance is always read 0.05 g higher than the actual value.
2. The side product that formed is Mg_3N_2 .
3. If the magnesium oxide is not dried completely the reported value of (Mg to O) mole ratio will decrease.
4. (Mg to O) mole ratio will increase if air is not sufficient to react with all the magnesium.
5. (Mg to O) mole ratio will decrease if a rapid oxidation is occur and some of the magnesium is lost.

Ans: 5

Q7) A compound of iridium, Ir (M.Wt = 192.2 g/mol), and oxygen, O (M.Wt = 16 g/mol), was produced in a lab by heating iridium in a crucible, the data was collected:

Mass of crucible 38.26 g

Mass of crucible and iridium 39.63 g

Mass of crucible and iridium oxide 39.74 g

What is the empirical formula of this compound?



Sol $\rightarrow$

$$\text{Mass Ir} = 39.63 \text{ g} - 38.26 \text{ g} = 1.37 \text{ g}$$

$$\text{Mass iridium oxide} = 39.74 \text{ g} - 38.26 = 1.48 \text{ g}$$


$$\text{Mass O} = 1.48 - 1.37 = 0.11 \text{ g}$$

$$\begin{array}{rcl} \text{Ir} & : & \text{O} \\ \frac{1.37 \text{ g}}{192.2} & & \frac{0.11 \text{ g}}{16} \\ \frac{7.12 \times 10^{-3}}{6.87 \times 10^{-3}} & : & \frac{6.87 \times 10^{-3}}{6.87 \times 10^{-3}} \end{array}$$

Ans: 2



Q8) A 0.1000 g sample containing C (FM=12), H (FM=1) and O (FM=16) only, and produced 0.1910 g of CO_2 and 0.1172 g of H_2O , what is the empirical formula of the compound?



Sol →

$$\text{Mass C} \rightarrow 0.1910 \text{ g CO}_2 \times \frac{1 \text{ mol CO}_2}{44 \text{ g CO}_2} \times \frac{1 \text{ mol C}}{1 \text{ mol CO}_2} \times \frac{12 \text{ g C}}{1 \text{ mol C}}$$

$$= 4.34 \times 10^{-3} \text{ mol C} = 0.052 \text{ g C}$$

$$\text{Mass H} \rightarrow 0.1172 \text{ g H}_2\text{O} \times \frac{1 \text{ mol H}_2\text{O}}{18 \text{ g H}_2\text{O}} \times \frac{2 \text{ mol H}}{1 \text{ mol H}_2\text{O}} \times \frac{1 \text{ g H}}{1 \text{ mol H}}$$

$$= 0.013 \text{ mol H} = 0.013 \text{ g H}$$

$$\text{Mass O} \rightarrow 0.100 - (0.052 + 0.013) = 0.035 \text{ g O} = 2.1875 \times 10^{-3} \text{ mol O}$$

C	:	H	:	O
$\frac{4.34 \times 10^{-3}}{2.1875 \times 10^{-3}}$		$\frac{0.013}{2.1875 \times 10^{-3}}$		$\frac{2.1875 \times 10^{-3}}{2.1875 \times 10^{-3}}$

2	:	6	:	1
---	---	---	---	---



Ans: 4

For exp 1

Q9) Which of the following statement is not correct?

1. Open Sandals, short skirts and shorts are not allowed in the lab.
2. When dealing with Flammable material, don't heat with direct flame.
3. You should return excess chemicals to their reagent bottles
4. Never taste or smell chemicals or solutions in the lab.
5. Laboratory work can't be started as soon as you enter the lab, unless the teacher is present.

Ans: 3

Q10) Given the equation $A + 3B \rightarrow C + D$ by reacting 1 moles of A with 2 moles of B, which of the following is true?

1. A is limiting reactant because of it's higher molar mass.
2. B is limiting reactant because of it's higher molar mass
3. A is limiting reactant because you have fewer moles of A than B
4. B is limiting reactant because you have fewer moles of A than B
5. B is limiting reactant because you need 3 moles of B and you have 2.

Sol $\rightarrow \frac{1A}{1} = 1A$ $\frac{2B}{3} = \underline{0.67B}$ B is LR

coefficient

Ans: 5

Q11) If 3.28 g unknown mixture containing $Na_3PO_4 \cdot 12H_2O$ (M.Wt = 380.2 g/mol) and $BaCl_2 \cdot 2H_2O$ (M.Wt = 244.2 g/mol) is dissolved in distilled water, The mass of $Ba_3(PO_4)_2$ (M.Wt = 602.2 g/mol) precipitate is 1.75 g. Calculate the % of $Na_3PO_4 \cdot 12H_2O$ in the mixture if the $BaCl_2$ is the limiting reactant.

1. 44.19%
2. 35.09%
3. 65.58%
4. 57.42%
5. 75.00%

Sol $\rightarrow \frac{3Ba^{+2} + 2PO_4^{-3}}{L.R} \rightarrow Ba_3(PO_4)_2$

$$1.75 \text{ g } Ba_3(PO_4)_2 \times \frac{1 \text{ mol } Ba_3(PO_4)_2}{602.2 \text{ g } Ba_3(PO_4)_2} \times \frac{3 \text{ mol } Ba^{+2}}{1 \text{ mol } Ba_3(PO_4)_2}$$

$$\times \frac{244.2 \text{ g } Ba^{+2}}{1 \text{ mol } Ba^{+2}} = 2.128 \text{ g } Ba^{+2}$$


$$\text{Mass } PO_4^{-3} = 3.28 - 2.128 = 1.15 \text{ g}$$

$$\% Na_3PO_4 \cdot 12H_2O = \frac{1.15}{3.28} \times 100 \% = 35.09\%$$

Ans: 2

of $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ (M.Wt = 380.2 g/mole) and $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ (M.Wt = 244.2 g/mole) is dissolved in water, A ppt. of $\text{Ba}_3(\text{PO}_4)_2$ (M.Wt = 602.2 g/mole) was produced with a mass $0.2x$ of the starting materials. Then the Percentage yield of $\text{Ba}_3(\text{PO}_4)_2$ is:-

- 1) 37.9% 2) 63.1% 3) 88.3% 4) 25.3%



$$\% \text{ yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100\%$$

$$\text{mass Ba}^{+2} = \text{mass PO}_4^{-3} = x \text{ g}$$

II Find the L.R

$$\text{Ba}^{+2} \rightarrow \frac{x \text{ g}}{244.2 \text{ g/mole}} = \frac{x}{244.2} \text{ mole} / \underset{\substack{\uparrow \\ \text{coefficient}}}{3} = \frac{x}{732.6}$$

$$\text{PO}_4^{-3} \rightarrow \frac{x \text{ g}}{380.2 \text{ g/mole}} = \frac{x}{380.2} \text{ mole} / 2 = \frac{x}{760.4} \text{ // smaller than}$$

So PO_4^{-3} is L.R

$$\frac{x}{380.2} \text{ mole PO}_4^{-3} \times \frac{1 \text{ mole Ba}_3(\text{PO}_4)_2}{2 \text{ mole PO}_4^{-3}} \times \frac{602.2 \text{ g Ba}_3(\text{PO}_4)_2}{1 \text{ mol Ba}_3(\text{PO}_4)_2}$$

$$= 0.791 x \text{ g} \equiv \text{Theoretical yield}$$

$$\% \text{ yield} = \frac{0.2x}{0.791x} \times 100\% = 25.3\%$$

Ans: 4

Q13) In the limiting reactant of salt mixture exp. to test for Limiting Ba^{+2} ion:-

Ans: 1

1. Add few drops of Barium ion (0.5 M) to the supernatant and ppt. will appear
2. Heat the mixture solution in the water bath for 30 minutes.
3. Add few drops of phosphate ion (0.5 M) to the supernatant and ppt. will appear
4. Add few drops of Barium ion (0.5 M) to the supernatant and ppt. will not appear
5. none of the above.

The Sulfate ion can be detected by:-

1. Adding BaCl_2 Solution in acidic media and a white ppt. will appear.
2. Adding BaCl_2 Solution in basic media and a white ppt. will appear.
3. Adding HCl Solution, a gas will change the wet red litmus to blue.
4. Adding NaOH Solution, a gas will change the wet blue litmus to red.
5. Adding NaOH Solution, ammonia smell can be detected.

Ans: 1

Q15) The Cl^- can be detected by:-

1. Sodium oxalate
2. Potassium thiocyanate
3. Silver nitrate + acid
4. Barium chloride + acid
5. Hydrochloric acid.

Ans: 3

Q16) When an unknown react with sodium ^{NaOH} hydroxide solution, it evolved a gas which convert the wet red litmus paper to blue. The resulted aqueous layer from the previous reaction was ~~the~~ treated with hydrochloric acid solution and carbon dioxide ^{H_2CO_3} evolved immediately as a result of reaction, The unknown is:-

1. CaCl_2
2. $\text{Ca}(\text{HCO}_3)_2$
3. NH_4Cl
4. NH_4HCO_3

Ans: 4

Q17) An unknown salt give a gas that convert the litmus Paper from red to blue when deteteel with sodium hydroxide and a pale yellow precipitate when reacted with silver nitrate in acidic media. The formula of the salt is:-

1. CaBr_2
2. $\text{Fe}_2(\text{SO}_4)_3$
3. NH_4HCO_3
4. FeBr_3
5. NH_4Br

Ans: 5

Q18) The iron (III) ion can be detected by:-

Ans: 2

1. Adding BaCl_2 solution, in acidic media and a white ppt. will appear.
2. Adding KSCN solution, and a red color will appear.
3. Adding HCl solution, a gas will change the wet red litmus paper to blue.
4. Adding NaOH solution, a gas will change the wet blue litmus paper to red.
5. Adding KSCN solution, and a white color will appear.

Q19) which of the following pair of liquids are miscible?

1. Polar + Salt
2. non-polar + non-polar
3. a and b will be miscible
4. Non-polar + Polar
5. Non-polar + Salt.

Ans: 3

Q20) which of the following statement is correct?

1. The boiling point is the temperature at which the vapor pressure of the liquid is higher than the atmospheric pressure
2. If the boiling point is recorded while the bubble escaping from the capillary tube (and the heat is removed), the recorded Boiling point will be too low
3. The boiling point of the substance increase as the intermolecular forces between molecules decrease.
4. If the boiling point is recorded after the liquid enters the capillary tube (after the heat is removed), the recorded boiling point will be too high.
5. As the temperature increases the vapor pressure of the liquid increases.

* Relations →

Ans: 5

- | | |
|----------------------------|------------------|
| ① Inter molecular Forces ↑ | vapor pressure ↓ |
| ② Temperature ↑ | vapor pressure ↑ |
| ③ Boiling point ↑ | vapor pressure ↓ |
- And So on ---

Q21) If you need 10 ml pipette to weight 10 ml of three unknowns liquid substances A, B and C. you find that the weight of the 10 ml of each substance is the following
 $A = 9.2 \text{ g}$, $B = 9.0 \text{ g}$, $C = 8.9 \text{ g}$. The order of density decreasing of these liquids is:-

1. $C > A > B$
2. $B > C > A$
3. $A > B > C$
4. $A > C > B$

Ans: 3

Q22)

Page 20

A student's liquid unknown boils at approximately 79°C , is insoluble in water but soluble in cyclohexane and ethanol. The mass of 2ml of the unknown = 1.75 g, from the table below the unknown is:

Compound	Density g/ml	Boiling Point ($^{\circ}\text{C}$)	Solubility		
			S: Soluble, i: insoluble		
			H_2O	C_6H_{12}	$\text{C}_2\text{H}_5\text{OH}$
X	0.79	68	S	S	i
Y	0.90	60	S	S	S
Z	0.59	78	i	S	S
R	0.89	80	i	S	S
Q	0.89	79	i	S	i
P	0.79	59	S	S	S

1. X

2. Z

3. Q

4. R

~~Density~~ Density = $\frac{m}{V} = \frac{1.75}{2} = 0.88 \text{ g/ml}$

Ans: 4

exp 4

Q23) The following results were obtained for an unknown containing a cation and an anion.

Cation test	Sodium oxalate	Potassium thiocyanate	(with litmus) Paper test Sodium hydroxide
	-ve	+ve	-ve
Anion test	Barium chloride (in acidic medium)	Silver nitrate (in acidic medium)	Hydrochloric acid (immediate reaction)
	+ve	-ve	-ve

1. FeCl_3 2. $(\text{NH}_4)_2\text{SO}_4$ 3. $\text{Fe}_2(\text{SO}_4)_3$ 4. NH_4Cl

Ans: 3

Exp 6:- Molar Mass of a volatile Liquid.

⇒ In this exp. we will compute the Molar Mass using:-

$$PV = nRT$$

$$PV = \frac{\text{Mass}}{M.M} RT \Rightarrow P = \text{atmospheric pressure (atm)} \Rightarrow 1 \text{ atm} = 760 \text{ mmHg} \\ = 760 \text{ torr} \\ = 101.325 \text{ Kpa}$$

$$\text{المذيب} \leftarrow V = \text{Volume of Vapor (L)} \Rightarrow 1 \text{ L} = 1000 \text{ mL}$$

$$n = \text{number of moles} = \frac{\text{mass (g)} \rightarrow \text{المذاب}}{M.M \text{ (g/mol)}}$$

$$R = \text{gas constant} = 0.0821 [\text{atm} \cdot \text{L} / \text{K} \cdot \text{mol}]$$

$$T = \text{temperature of boiling water K}$$

$$K^{\circ} = C^{\circ} + 273$$

1) Calculation part:-

تعويض مباشر في القانون مع مراعاة استخدام الوحدة المذكورة في القانون أعلاه.

ExA1) A flask weighs 40.1305g when clean, dry, evacuated, 138.2410g when filled with water (density = 0.9970 g/mL) and 40.2487g when filled with a gaseous substance at 470.4 torr and 96°C. What is the molar mass (g/mol) of the gas? [R = 0.0821 L.atm/mol.k]

$$\text{Solution} \rightarrow PV = \frac{\text{mass}}{M.M} RT$$

$$P(\text{atm}) = 470.4 \text{ torr} \times \frac{1 \text{ atm}}{760 \text{ torr}} = 0.61895 \text{ atm}$$

$$V_{\text{(L)}} = \frac{\text{Mass water}}{\text{density}} \Rightarrow \text{Mass water} = \text{Mass flask filled with water} \\ - \text{Mass empty flask} \\ = 138.2410 - 40.1305 \\ = 98.1105 \text{ g}$$

$$V = \frac{98.1105 \text{ g}}{0.9970 \text{ g/mL}} = 98.4057 \text{ mL} = 0.0984057 \text{ L}$$

$$\text{Mass (g)} = \text{Mass Flusk with substance} - \text{Mass empty flusk}$$

(gaseous substance)

$$= 40.2487 - 40.1305 = 0.1182 \text{ g.}$$

$$R = 0.0821$$

$$T = 96 + 273 = 369 \text{ K}$$

⇒ Now:-

$$PV = \frac{\text{Mass}}{M.M} RT \Rightarrow M.M = \frac{\text{Mass} \cdot R \cdot T}{PV}$$

$$= \frac{(0.1182)(0.0821)(369)}{(0.61895)(0.0984057)}$$

$$= 58.79 \text{ g/mole}$$

* We can rearrange the equation to calculate density:-

$$PV = \frac{\text{Mass}}{M.M} RT \Rightarrow \frac{P(M.M)}{RT} = \frac{\text{Mass}}{\text{Volume}} \Rightarrow D = \frac{P(M.M)}{RT}$$

D: Density (g/L)

PRE LAB:-

1) For which of the following compounds can we determine its molar mass using the method described in this experiment? Give reasons. Benzene (b.p. 78°C), Glycerol (b.p. 180°C)

Solution → دائماً نختار المادة التي درجة غليانها أقل من الماء فشان تبخر قبل ما يغلي الماء.

Benzene (78°C) because its boiling pt lower than water.b.p.

2) A cylinder contains compressed hydrogen gas and the mass of the hydrogen is 20 g, what mass of oxygen would be contained in an identical cylinder at the same temperature and pressure?

Solution →

since the cylinder is identical $\Rightarrow V_O = V_H$

using avogadro's principle: $\frac{n_H}{V_H} = \frac{n_O}{V_O} \Rightarrow n_H = n_O$

$$n_H = 20 \text{ g} \times \frac{1 \text{ mol}}{1 \text{ g}} = 20 \text{ mol} = n_O$$

$$\text{Mass O} = 20 \text{ mol} \times \frac{16 \text{ g}}{1 \text{ mol}} = 320 \text{ g}$$

2) Procedure part :-

- why should the Erlenmeyer flask be dry?
→ to avoid changing the mass of the substance.
- why we should make a small hole through the aluminum foils? to avoid the explosion of the flask.

- Does it effect if we change the quantity of the water? No, M.M is independent on the quantity of the water

يعني لو مكاني كان عندي 5 mL من الماء و صارو 10 mL بيختلف اشي؟
الجواب لا، لأنني ما باخذ الحجم من الماء الموجود، ببساطة ليتبخر و بس
يخلص تبخر باخذ حجم البخار وليس السائل.

- Why we are putting a few boiling stones (boiling chips) in the liquid? to decrease the bubbles of the boiling so we can avoid explosion.

- why should we heat the beaker slowly?
To avoid fast evaporation
مستأن ما تخلص الماء بسرعة
وبالتالي ما تتم عملية ال Vaporization كاملة.

- why should we dry the outside of the aluminum foil completely, after finish heating?
To have an accurate reading of the mass.

Questions:-

Q₁) Describe the effect of the following factors (whether increase, decrease, or has no effect) on the calculated molar mass of the volatile liquid.

a. IF the flask was not dried well before the weighing.
 a. decrease ~~b. increase~~ c. No effect.

$$M.M = \frac{\text{Mass}}{PV} \cdot RT \Rightarrow \text{was not dried well} \Rightarrow \text{mass} \uparrow \therefore M.M \uparrow$$

b. IF the volume of the flask is bigger than the recorded volume.

يعني V المحسوبة أقل.

$$V \downarrow, M.M \uparrow$$

~~a. increase~~ b. decrease c. No effect.

c. IF the temperature of the boiling water was mistakenly less than the true value.

a. increase ~~b. decrease~~ c. No effect.

$$T \downarrow, M.M \downarrow$$

d. IF the density of the volatile liquid was mistakenly greater than the true value.

$$M.M = \frac{D}{P} RT, D \uparrow, M.M \uparrow$$

~~a. increase~~ b. decrease c. No effect.

Q₂) what is the mass of vapor of volatile liquid (M.Wt. = 85 g/mol)

which completely fill a 184 ml flask at 94.0 °C and 675.05 torr?

a. 0.211 g b. 0.632 g ~~c. 0.461 g~~ d) 0.344 g.

Solution $\Rightarrow$

$$\text{Mass} = \frac{PV M.M}{RT} = \frac{\left(\frac{675.05}{760}\right)(0.184)(85)}{0.0821(94+273)} = 0.461 \text{ g}$$

Exp 8:- Colligative properties:- Molar Mass Determination.

Objective:- To determine the molar mass of a non-volatile, non electrolyte by observing the difference between the freezing points of a solvent and a solution.

non-volatile $\rightarrow$ غير متطايرة, non-electrolyte $\rightarrow$ غير متفككة

$\rightarrow$ when we add a non-volatile solute to a solvent it changes the physical properties of the solvent:-

- 1) Freezing pt (F.P) $\Rightarrow$ decreasing (Freezing pt depression) ^{الانخفاض}
- 2) Boiling pt (B.P) $\Rightarrow$ increasing (Boiling pt elevation) ^{ارتفاع}
- 3) Vapor pressure (V.P) $\Rightarrow$ decreasing (Vapor pressure lowering) ^{انخفاض}

$\Rightarrow$ Freezing pt depression, Boiling pt elevation & Vapor pressure lowering are called Colligative properties.

$\Rightarrow$ And they are governed by number, rather than type
يعني ذاتها Quantitative بتحدد على الكمية وليس على النوع.

$$\Rightarrow \Delta T_f = K_f m = K_f \cdot \frac{(\text{g mass solute})}{\text{M.M. Solute} \cdot \text{kg Solvent}} = \frac{K_f * \text{وزن المذاب بالجرافان}}{\text{وزن المذيب بالـ Kg} \times \text{مولر ماس المذاب}}$$

Freezing pt depression $\rightarrow$ نصاب التجربة رج تركيز على
Boiling pt elevation. $\rightarrow$ أكثر من

$\Rightarrow K_f$ and K_b are the molar Freezing pt and boiling pt constants for the solvent. (Given)

$$\Rightarrow m; \text{ molality} = \frac{\text{moles solute (mole)}}{\text{Mass Solvent (Kg)}}$$

* IF we have an electrolyte solute $\rightarrow$ مذاب يتفكك
we will have a Van't Hoff Factor (i) $\rightarrow$ معامل التفكك.
which leads to:- ^{For non-electrolyte i=1.}

$$\Delta T_f = i K_f m \Rightarrow \text{بمعنى لو كانت المادة متفككة يعني إنو الفرق في درجة الحرارة (الانخفاض) يزداد.}$$

* لازم نغير بين درجة الحرارة T_f وبين الانخفاض في درجة الحرارة ΔT_f
 $\Delta T_f = T_i - T_f \Rightarrow$ ازدياد الانخفاض في درجة الحرارة ΔT_f
يعني نقصان في درجة الحرارة النهائية $\downarrow T_f$

Pre Lab:

- 1) Students prepared two cyclohexane solutions having the same mass of solute. However student 1 used 13g of cyclohexane, student 2 used 15g. which student will observe the larger freezing pt. change? Explain

$$\Delta T_f = K_f \frac{\text{Mass (solute)} \rightarrow \text{كتلة المذاب}}{M.M \text{ (Mass Solvent)} \rightarrow \text{كتلة المذيب}} \Delta T_f$$

$$\Delta T_{f \text{ student 1}} > \Delta T_{f \text{ student 2}}$$

Since Mass solvent 1 < Mass solvent 2.




- 2) A 0.597 g sample of a non-electrolyte dissolves in 20.0 g of cyclohexane. The freezing point depression is 3.62°C . What is the molar mass of the non-electrolyte? (K_f for cyclohexane is $20.0^\circ\text{C kg/mol}$).

Solution $\Rightarrow$

$$\Delta T_f = K_f \frac{\text{mass solute (g)}}{M.M \text{ solute} \times \text{Mass Solvent (kg)}}$$


$$M.M = \frac{20 \times 0.597}{3.62 (20 \times 10^{-3})} = 164.92 \text{ g/mol}$$

i For some solutes:-



Procedures part:-

- Why should we keep moving the solution when it freezes?
To avoid super cooling. عشان مانتفضح درجة الحرارة كثير
- Cooling curve :- Freezing pt depression as a function of time.
- The temperature remains constant at the freezing pt. untill the Freezing is almost complete.



Post Lab:-

- 1) If the solution's Freezing pt is erroneously read 0.2°C lower than it should be, will the unknown's calculated molar mass be too high or too low? Explain!

Solution $\Rightarrow \Delta T_F = T_{\text{solvent}} - T_{\text{solution}} = \frac{K_F \text{ mass solute}}{M.M \text{ solute} \times \text{mass solvent}}$

$$T_{\text{solution}} \downarrow \Rightarrow \Delta T_f \uparrow$$

$\Delta T_f \uparrow \Rightarrow M.M \downarrow \therefore$ Molar Mass will decrease.

- 2) How will the freezing pt. change of cyclohexane be affected (compared with the freezing pt. change by a non-volatile, non-associating and non-dissociating solute) by:-

الوجه، عندي
ارتباط معي

4. معرفة فلك $i=1$

- (a) A non-volatile solute that dissociates? Explain!

عَاجِلُو رَح سَيِّفُكَلْ إِذَا رَح يَكُون عِنْدِي مَعَامِل تَفَكَلْ نَا

Solution $\rightarrow \Delta T_f = i K_f m \rightarrow i \uparrow \rightarrow \Delta T_f$ increase.

- (b) Two solutes that react according to the equation.

$A+B \rightarrow C?$ Explain!

1 1

 $i = 2$ *i*

هون چهار ترابط كانت $i=2$ چهار $i=1 \Rightarrow$

$$i=2 \quad i=1 \Rightarrow$$
$$\Delta T_f = i k_f m \Rightarrow i \downarrow \Rightarrow \Delta T_f \text{ decrease.}$$

3) IF some solute adheres to the test tube's wall in part B.1, is the freezing point change greater or less than it should be? Explain!

في التجربة لما صطينا المذاب مشان يذوب ونقيس درجة التجمد بيسألني إذا التصق جزء من المذاب سؤرح ياتر على مقدار الانخفاض؟

Solution $\rightarrow \Delta T_f = \frac{i K_f (\text{Mass solute})}{M.M (\text{Mass solvent})} \Rightarrow$ Mass solute $\downarrow$
 $\therefore \Delta T_f$ will decrease.
 (Be less than it should be).

* ما نسي نفتر بين ΔT_f و T_f .

Questions:-

Which statements of the following statements are T and which of them are F?

1. (T) IF the test tube contains an insoluble impurity, then the calculated molar mass will be no effected.

$$\Delta T_f = \frac{i K_f \times \text{Mass solute}}{M.M \times \text{Mass solution}}$$

لأنه هون بس بحسب الكتلة اللي ذابت اللي ما ذابت ما الي فيها.

2. (T) IF the thermometer reading is always 1.5 °C higher than the correct temperature, the calculated molar mass will be not effected.

له لانه رح تقرأ 1.5 أعلى لما أقيس المذيب الخالي و 1.5 أعلى لما أقيس المحلول لما أخذ القزعة بدرجة مع بعض المشكلة لو اختلفت القراءة عند وحدة منهم بس.

3. (F) The Freezing pt of the solution had been incorrectly read 0.6 °C higher than the true Freezing pt, the calculated Molar Mass will be lower than actual.

$$\Delta T_f = T_{\text{solvent}} - T_{\text{solution}}$$

$$T_{\text{solution}} \uparrow \Rightarrow \Delta T_f \downarrow$$

$$\Delta T_f = \frac{K_f \cdot \text{mass(g)}}{M.M \cdot \text{mass(kg)}}$$

$$\Delta T_f \downarrow \Rightarrow M.M \uparrow$$

4) (F)

$$\Delta T_f = i K_f \frac{\text{moles}}{\text{Mass}_{\text{solvent}}}$$

↖ ← ← ← ↘

$$\begin{matrix} i_{\text{NaCl}} = 2 \\ i_{\text{C}_{10}\text{H}_8} = 1 \end{matrix} \rightarrow \Delta T_f \text{NaCl} > \Delta T_f \text{C}_{10}\text{H}_8$$

The Freezing pt. depression of 0.20 mole of NaCl in 10 g of water is lower than the Freezing pt depression of 0.20 mole C_{10}H_8 in 10.0 g of water.

Q₂) A solution of 3.33 g of unknown in 50 g of water Freezes at -0.773°C , What is the molecular weight of the unknown? $K_f = 1.86^\circ\text{C/molal}$
M.M?

$$T_f \text{ solvent} = T_f \text{ water} = 0$$

$$\Delta T_f = 0 - (-0.773) = i \frac{K_f (\text{mass solute})}{\text{M.M}_{\text{solute}} * \text{Mass solvent}}$$

$$\text{M.M} = \frac{1 (1.86) (3.33)}{0.773 (0.05)} = 160$$

- a) 120
- ✓ b) 160
- c) 80
- d) 100

→ ΔT_f is always +ve

وإذا ما كانت موجبة بإختصار
التي المثلثة

Objective: To measure heat of reaction.

Calorimetry → It's the measurement of heat change.

Calorimeter → It's a device that used to measure the heat of the (reaction) (rxn).

↳ There are several kinds of "heats of reactions" such that → 1) heat of solution
2) heat of neutralization.

$$\Delta H_{\text{reaction}} = \Delta H_{\text{solution}} + \Delta H_{\text{neutralization}}$$

$$\Delta H_{\text{rxn}} = \Delta H_{\text{sol}} + \Delta H_n$$

• heat of solution → heat flows during a process of solution.
OR → amount of heat required all released to form a solution.

heat of neutralization → amount of heat required all released to make a neutralization.

heat of reaction → It's the total heat $\Delta H_n + \Delta H_{\text{sol}}$.

$\Delta H_{\text{rxn}} \rightarrow$ يمكن حسابها عملياً $\Delta H_n \rightarrow$ لا يمكن حسابها عملياً

$\Delta H_{\text{sol}} \rightarrow$ يمكن حسابها عملياً

* توضيح للي بصير فوق :-




أنا بدي أعمل تفاعل التفاعل اللي هو


وبدي أحسب الحرارة الناتجة من هذا التفاعل بين ما بقدر أحسبها مباشرة.

كيف نحسبها ؟

EXAMPLE)




2) $\Rightarrow$
يكون



1) بحسب الحرارة الناتجة من ذوبان المحلول

محلول الناتج

2) بحسب الحرارة الناتجة من تفاعل مادة مع محلول الناتج



$\Delta H_{rxn} \rightarrow$ حرارة المحلول + حرارة تفاعل التبادل

$$\Delta H_{rxn} > \Delta H_n, \rightarrow \Delta H_{sol}$$

$$\Delta H_n = \Delta H_{rxn} - \Delta H_{sol}$$

$$\Delta H = (-) \text{Mass of Solution (g)} \times \text{specific heat } \left(\frac{J}{g \cdot ^\circ C} \right) \times \Delta T (^\circ C)$$

Solute + Solvent

$$\Delta H = () J$$

or $\Delta H = (-) \frac{\text{mass solution}}{\text{moles of solute}} \times \text{specific heat} \times \Delta T (^\circ C) \times 10^3$

$$\Delta H = () \frac{\text{mass}}{\text{M.M}} KJ/mol$$

$\Delta H \rightarrow -ve (T_f > T_i) \rightarrow$ exothermic (طارد للحرارة) $\rightarrow$ Flows out of the system. (بالنوع)
 $\Delta H \rightarrow +ve (T_f < T_i) \rightarrow$ endothermic (ماص للحرارة) $\rightarrow$ Flows into the system.

* Specific heat: Amount of heat required to raise the temperature of 1 g mass by $1^\circ C$. (بالمقدار)

* $\Delta T \rightarrow$ Change in temperature.

Example 1) A 2.00 g sample of solid CsOH is dissolved in 200.0 mL of water in a calorimeter. The temperature of the water was raised from 22.3 to $23.4^\circ C$, calculate the heat of the solution in KJ/mol . (assume the specific heat of the solution to be $4.184 J/g^\circ C$ and the density of the solution to be $1 g/mL$)?

Solution $\rightarrow \Delta H = - \text{Mass}_{\text{solution}} \times \text{specific heat} \times \Delta T$

$$\Delta T = T_f - T_i = 23.4 - 22.3 = 1.1^\circ C$$

$$\text{specific heat} = 4.184$$

$$\text{mass solution} = \text{mass solute} + \text{mass solvent}$$

$$= 2g + 200g = 202g$$

$$\Delta H = -(202)(4.184)(1.1) = -930 J$$

in $KJ/mol \Rightarrow \Delta H = \frac{-930}{(2/149.9)} \times 10^{-3} = -70 KJ/mol$ ΔH_{sol}

mass CsOH/M.M

$$m = d \times V$$

$$= 1(200) = 200g$$

Example 2) A 2.00 g sample of solid CsOH reacted with 200.00 mL of aqueous solution of hydrochloric chloride (HCl) in a calorimeter, the temperature of the solution increased from 22.3 to 24.3 °C. calculate the heat of reaction in KJ/mol? (Assume the specific heat of the solution to be 4.184 J/g°C and the density of the solution to be 1.00 g/mL)

جول
در
درجة
C
80
ΔH_{rxn}

$$\begin{aligned} \text{Sol} \rightarrow \Delta H &= - \text{mass}_{\text{solution}} \times \text{specific heat} \times \Delta T \\ &= - (2 + 200) \times 4.184 \times (24.3 - 22.3) \\ &= -1700 \text{ J} \end{aligned}$$

$$\text{in KJ/mol} = \frac{-1700}{\left(\frac{2}{149.9} \right)} = -130 \text{ KJ/mol} \quad \Delta H_{\text{rxn}}$$

→ From the previous 2 examples:

$$\Delta H_u = \Delta H_{\text{rxn}} - \Delta H_{\text{sol}} = -130 - (-70) = -60 \text{ KJ/mol.}$$

Explo Electrochemistry

→ Any chemical rxn. involves the transfer of e^- s from one substance to another is an Oxidation-Reduction Rxn.

← التفاعلات التي يتم فيها انتقال الإلكترونات من مادة إلى أخرى تسمى تفاعلات التأكسد والإختزال.

↳ Oxidation → loss of e^- s

↳ Reduction → gain of e^- s.

↳ A galvanic cell exists when the Oxidation & reduction steps in the rxn takes place, so that e^- s transfer from the reducing agent (substance which is oxidized) to the oxidizing agent

↳ E_{cell} consists of ^{قطب}cathode, ^{مصعد}anode, ^{قنطرة ملحية}salt bridge & Voltmeter.

Cathode: electrode at which ^{reduction}~~oxidation~~ occurs (+ve electrode)


Anode: electrode at which oxidation occurs (-ve electrode)

↳ e^- s flow from anode to cathode.

← لما يصير عندي تفاعل تأكسد وإختزال رح تنتقل عندي الإلكترونات من القطب السالب إلى القطب الموجب.
المصعد إلى المربط.
العامل المؤكسد إلى العامل المؤكسد المختزل.

القطب يلي صار عليه التأكسد إلى القطب يلي صار عليه الإختزال.
داخل ما يسمى بالخلية الجلفانية.

أقل ٥٥ يعكس المعادلة ويغير الإشارة




هون دايقا بيحطيني جهد الإختزال
أنا بتطلع على الأيسر ويعكس إشارته
ومعادلته تكون هاد هو طرف التأكسد
والثاني يكون طرف الإختزال

الإلكترونات بالتوازي نصف الأكسدة




$\epsilon^0 = 0.76 V$ (Anode)



$\epsilon^0 = 0.34 V$ (cathode)


نصف الأكسدة
الإلكترونات بالتوازي



↳ ① The first half-reaction is the oxidation half cell

↳ ② The second half-reaction is the reduction half cell.

→ The galvanic cell:-



In the previous example.

$\epsilon_{Cu^{+2}/Cu}^0 > \epsilon_{Zn^{+2}/Zn}^0$


بما أن $\epsilon_{Cu^{+2}/Cu}^0$ أكبر من $\epsilon_{Zn^{+2}/Zn}^0$ ، إذن $Cu^{+2}/Cu \rightarrow$ reduction half
و $Zn/Zn^{+2} \rightarrow$ Oxidation half.

Also we should note:-

if $\epsilon_{cell}^0 > 0 \Rightarrow$ Rxn is spontaneous in the written direction

if $\epsilon_{cell}^0 < 0 \Rightarrow$ Rxn is non-spontaneous in the written direction

The previous cell rxn. can be represented by the following Line cell:-



→ The salt bridge is usually a tube that is filled with standard electrolyte soln. Such as KCl , KNO_3 --- etc and the purpose of it is to compensate the ions migration by providing the soln. that has a cation migration by cations.

← تعويض النقص في أيونات المحلول.

↳ In the previous example (E° value were taken from a standard potential table at standard conditions: $25^\circ C$, 1 atm & $1M$)

↳ E_{cell} at conditions rather than the standard conditions can be calculated by Nernst equation:-

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{2.303 RT}{n F} \ln Q$$

$$= E_{\text{cell}}^\circ - \frac{0.0592}{n} \log Q \quad (\text{at } 25^\circ C) \quad \leftarrow \text{هنا الذي نستخدمه}$$

($R = 8.314 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$, $F = \text{Faraday's Const} = 96500 \text{ C/mole } e^-$
 $n = \text{no of } e^- \text{ mole}$)

Q :- The product of molar concentration of products divided by the product of molar concentrations of reactions, if there was a gas so we use the partial pressure of that gas.

$$Q = \frac{[\text{محاصل ضرب تراكيز النواتج}]}{[\text{محاصل ضرب تراكيز المتفاعلات}]}$$

① لو كان عندي غاز P باخذ بدل من التركيز.

② المواد التي باخذها التراكيز بين
 أو الضغط

Exp7, Determination of the Molar Volume of Hydrogen gas.

$$\text{Molar Volume} = \bar{v} = \left(\frac{\text{Volume of gas (L)}}{\text{moles of gas (mole)}} \right) = \frac{V}{\text{moles}}$$

L/mole

الظروف
المعيارية

STP → $T = 0^\circ\text{C} = 273\text{ K}$, $P = 1\text{ atm} = 760\text{ torr}$
 $V = 22.4\text{ L} \rightarrow$ for water (only)

$\bar{v} = 22.4\text{ L/mole} \rightarrow$ STP

Gases Laws:-

1) Boyle's Law → $P_1 V_1 = P_2 V_2$
 $P \propto \frac{1}{V}$

(درجة الحرارة ثابتة)



2) Charles's Law → $\frac{V_1}{T_1} = \frac{V_2}{T_2}$

(الضغط ثابت)

$V \propto T$

3) The Combined gas Law →

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

لما بدى أعمل حساباتي باستخدام عملية البتر بوجود الماء راح أحسب الضغط الكلي يلي هو ضغط الماء + ضغط الغاز يلي بدى أجمعه
 $P_{\text{total}} = P_{\text{H}_2\text{O}} + P_{\text{gas}}$

بس أنا بحسباتي راح أستخدم ال P_{gas} مشان أطلع أي شيء، والعلاقة فيه اللي هو

$$P_{\text{gas}} = P_{\text{total}} - P_{\text{H}_2\text{O}}$$

PreLab:-

A student at the Hashemite university wants to determine experimentally the volume occupied by one mole of H_2 gas at STP. She reacts 0.1471 g of Zn with excess $HCl(aq)$ and collects 56.09 ml of gas over water at $22^\circ C$ and 757.8 torr. The vapor pressure of water at $22^\circ C$ is 19.8 torr.

1- Use data given above to calculate


(i) The volume occupied by one mole of dry H_2 at $22^\circ C$, 760 torr

(ii) The volume occupied by one mole of dry H_2 at STP

This is Molar Volume, $\bar{V}$

Sol $\rightarrow$ (i)

① حتى حسب $\bar{V} = \frac{V_{gas}}{n_{gas}}$ لازم جيب n من المعادلة $\leftarrow$ من قانون الغازات



$$0.1471 \text{ g Zn} \times \frac{1 \text{ mol Zn}}{65.4 \text{ g Zn}} \times \frac{1 \text{ mol H}_2}{1 \text{ mol Zn}} = 2.249 \times 10^{-3} \text{ mol H}_2$$

من المعادلة $\times$

من الجدول الدوري

$$V_{1H} = 0.0569 \text{ L} \quad T_1 = 22 + 273 = 295 \text{ K}$$

$$\frac{P_{1\text{gas}} V_{1\text{gas}}}{T_1} = \frac{P_{2\text{gas}} V_{2\text{gas}}}{T_2} \quad \leftarrow \text{هنا نستخدم قانون الغازات المثالية}$$

السؤال أعطاني الضغط البخاري للماء (Over water) المجموع فوق الماء.
بلي هو

$$P_{\text{total}} = P_{H_2O} + P_{H_2}$$

$$19.8 \text{ torr} \quad \leftarrow \text{عطاني}$$

$$P_{H_2} = 757.8 - 19.8 = 738 \text{ torr.}$$

هنا نستخدم قانون الغازات المثالية

$$\frac{P_{1H_2} V_{1H_2}}{T_1} = \frac{P_{2H_2} V_{2H_2}}{T_2}$$

$$P_{2H_2} = 760 \text{ torr}$$

$$T_2 = 295 \text{ K.}$$

$$\frac{738 (0.0569)}{295} = \frac{760 (V_2)}{295} \Rightarrow V_2 = 54.4 \text{ mL.}$$

$$\bar{V} = \frac{54.4 \times 10^{-3}}{2.249 \times 10^{-3}} = 24.188 \text{ L/mole}$$

Sol 2 (ii) →

$$P_{1H} = 738 \text{ torr}$$

$$V_{1H} = 0.0569 \text{ L}$$

$$T_1 = 295 \text{ K}$$

$$P_{2H} = 760 \text{ torr}$$

$$V_{2H} = ?$$

$$T_2 = 273$$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow V_2 = \frac{738 (0.0569) (273)}{295 (760)}$$

$$= 50.4 \times 10^{-3} \text{ L} = 50.4 \text{ mL}$$

$$\bar{V} = \frac{V}{n} = \frac{50.4}{2.249} = 22.4 \text{ L/mol}$$

↪ at STP

2) Name the gas Laws which you used in your calculations?

1- Boyle's law , 2- Charles's Law

3- The combined gas law.

Example) Sample of KClO_3 decomposed producing O_2 gas that collected over water, The volume of the gas is 0.25 L at 26°C and 765 mmHg as total pressure, How many grams of KClO_3 was decomposed? , M.M of $\text{KClO}_3 = 122.6$
 $P_{\text{H}_2\text{O}}$ at $26^\circ\text{C} = 25 \text{ mmHg}$

Sol $\rightarrow$ $P_{\text{O}_2} V_{\text{O}_2} = \frac{n_{\text{O}_2} RT}{1}$

$$P_{\text{O}_2} = P_{\text{total}} - P_{\text{H}_2\text{O}} = 765 - 25 = 740 \text{ mmHg} = \frac{740}{760} \text{ atm}$$

$$V_{\text{O}_2} = 0.25 \text{ L}$$

$$\frac{740}{760} (0.25) = n_{\text{O}_2} (0.0821) (26 + 273)$$

$$n_{\text{O}_2} = 9.916 \times 10^{-3} \text{ mol O}_2$$

$$9.916 \times 10^{-3} \text{ mol O}_2 \times \frac{2 \text{ mol O}}{1 \text{ mol O}_2} \times \frac{1 \text{ mol KClO}_3}{3 \text{ mol O}} \times \frac{122.6 \text{ g KClO}_3}{1 \text{ mol KClO}_3} = 0.81 \text{ g}$$



← المتجيب معادلة بس، المتجيب المعلومة ←

→ Test banks :-

Page 4)

Q₁) A solution of 1.25 g of erythritol in 50 g of water freezes at -0.773°C , What is the molecular weight of erythritol? $K_f = 1.86^{\circ}\text{C/molal}$

- 1) 120 2) 60 3) 80 4) 100 5) 160

$$\Delta T_f = K_f \cdot \frac{\text{mass Solute (g)}}{\text{M.M} \times \text{mass Solvent (kg)}} \quad \rightarrow \text{M.M} = \frac{1.86 (1.25)}{(0 + 0.773)(0.05)} = 60.15$$

Q₂) In which of the following cases, the calculated molar mass of a volatile liquid will be lower than the actual value :-

- 1) The measured volume of the vapor was mistakenly larger than the true value. $V \uparrow \text{ M.M} \downarrow$
- 2) The temperature used in the calculations was higher than the actual boiling pt. of water under lab. conditions. $T \uparrow \text{ M.M} \uparrow$
- 3) The flask was not dried well before weighing. $\text{mass} \uparrow \text{ M.M} \uparrow$
- 4) The measured atmospheric pressure was less than the actual pressure. $P \downarrow \text{ M.M} \uparrow$
- 5) The measured density of the vapor was more than the actual one. $D \uparrow \text{ M.M} \uparrow$

$$\text{M.M} = \frac{\text{mass} RT}{PV} = \frac{DRT}{P}$$







Q₃) A conical flask weighs 40.1305 g when clean, dry, evacuated, 138.2410 g when filled with water at 25 C and 40.2487 g when filled with a gaseous substance at 300 torr and 96 C. What is the molar mass (g/mol) of the gas?

- 1) 92.2 2) 63.2 3) 27.4 4) 35.7 5) 42.5


mass solute = $40.2487 - 40.1305 = 0.1182 \text{ g}$
 Volume water = $138.2410 - 40.1305 = 98.1105 \text{ ml}$ / $P = \frac{300}{760}$
 Assume density 1g/ml
 $\text{M.M} = \frac{\text{mass} RT}{PV} = 92.2$

Q4) Among the following, which element can reduce


له صار اختزال
سؤال العنصر الذي تأكسد؟



هون بدى ادور على العنصر اللي جرد اختزاله اقل من جرد اختزال X
لانه مع الاختزال الاكبره اللي جرد تكون ذهب الاختزال والاكبره
رج تكون ذهب التاكسد بس بدى انتبه



Q5) Calculate the amount of heat liberated by dissolving 0.03 mol of $AlCl_3$ (M.wt = 133.33 g/mol) in 100 g water? If you know that the heat of solution is -321 kJ/mol



ΔH بوحدة kJ/mol

سؤال مباشر اعملك

حولها ل ΔH (kJ)

$$\Delta H = \frac{\Delta H}{\text{kJ/mol}} \times X \text{ mol}$$

$$= -321 \times 10^3 \frac{\text{kJ}}{\text{mol}} \times 0.03 \text{ mol} = -9.63 \text{ kJ}$$

Q6) In the Freezing point depression experiment, which of the following factors will increase the calculated molar mass of the solute:-

- 1) The solute has been totally dissolved in the solvent. $\rightarrow$ No effect.
- 2) The solution Freezing pt was erroneously lower than it should be. $\Delta T_f = T_{\text{solvent}} - T_{\text{solution}} \rightarrow \Delta T_f \uparrow, \text{M.M} \downarrow$
- 3) Some of the solvent evaporated during the experiment. $\text{mass solvent} \downarrow, \text{M.M} \uparrow$
- 4) Some solute adheres to the test tube. $\text{Mass} \downarrow, \text{M.M} \downarrow$
- 5) None of the above.

$$\text{M.M} = \frac{K_f \cdot \text{mass solute}}{\text{Mass solvent} \cdot \Delta T_f}$$

Q7) In the concentration cell, which of the following statements is not correct?

- 1) Electrons will flow from the lower concentration to the higher one.
- 2) The anode is the lower concentration electrode.
- 3) The standard potential is one. $\rightarrow$ Zero
- 4) The two half cells contain the same solution that differ in the concentrations only.
- 5) The cathode is the higher concentration electrode.

* هاد السؤال مهم ركزو على الخيارات الصحيحة 1, 2, 4, 5

Q8) A 1g sample of solid (NaOH, Mwt. = 40 g/mol) is dissolved in 100 mL of water in a Calorimeter the temperature of water was raised from 22.3°C to 23.8°C, Calculate the heat of the solution in kJ/mol ($S_p = 4.184 \text{ J/g}^\circ\text{C}$, and density of the solution 1g/mL)

- 1) -25.46 2) -9.19 3) -12.07 4) -17.97 5) -35.5

$$\Delta H = \frac{-\text{Mass solution} \times S_p \times \Delta T}{\text{Moles solute}} = \frac{-(100+1) \times 4.184 \times (23.8-22.3)}{1/40} = -25.46$$

Q9) If the Freezing point of the solution had been incorrectly read 0.3°C higher than its true Freezing point and the Freezing point of the pure solvent was correctly read, the effect on the calculated molar mass of the unknown is

- 1) Too high because change in temperature direct Proportional to molar mass.
- 2) Too Low because change in temperature direct Proportional to molar mass.
- 3) Too low because change in temperature inversly Proportional to molar mass
- 4) Too high because change in temperature inversly Proportional to molar mass.
- 5) No effect because the temperature does not change Significantly

$$\Delta T_f = T_{\text{solvent}} - T_{\text{solution}} \uparrow$$

$\Delta T_f \downarrow$

$$M.M = \frac{\text{mass}_{\text{solute}} \cdot K_f}{\Delta T_f \cdot \text{Mass}_{\text{solvent}}}$$

inversly

$$\Delta T_f \downarrow, M.M \uparrow$$

Q10) Among the Following, the weakest oxidizing agent is

- 1) Cu^{+2} (+0.34V)
- 2) Al (-0.66V)
- 3) Pb^{+2} (-0.13V)
- 4) Zn^{+2} (-0.76V)
- 5) Cr^{+2} (-1.45V)

مستقل اقوان
جهد اقوان
مستقل اقوان
جهد اقوان

- Cu^{+2} (+0.34V)
- Al^{+3} (+0.66V)
- Pb^{+2} (-0.13V)
- Zn^{+2} (-0.76V)
- Cr^{+2} (-1.45V)

افضل
عالم مؤيد
اقل
افضل

The Sulfate ion can be detected by:-

1. Adding BaCl_2 Solution in acidic media and a white ppt. will appear.
2. Adding BaCl_2 Solution in basic media and a white ppt. will appear.
3. Adding HCl Solution, a gas will change the wet red litmus to blue.
4. Adding NaOH Solution, a gas will change the wet blue litmus to red.
5. Adding NaOH Solution, ammonia smell can be detected.

Ans: 1

Q15) The Cl^- can be detected by:-

1. Sodium oxalate
2. Potassium thiocyanate
3. Silver nitrate + acid
4. Barium chloride + acid
5. Hydrochloric acid.

Ans: 3

Q16) When an unknown react with sodium ^{NaOH} hydroxide solution, it evolved a gas which convert the wet red litmus paper to blue. The resulted aqueous layer from the previous reaction was ~~the~~ treated with hydrochloric acid solution and carbon dioxide ^{H_2CO_3} evolved immediately as a result of reaction, The unknown is:-

1. CaCl_2
2. $\text{Ca}(\text{HCO}_3)_2$
3. NH_4Cl
4. NH_4HCO_3

Ans: 4

Q17) An unknown salt give a gas that convert the litmus Paper from red to blue when detected with sodium hydroxide and a pale yellow precipitate when reacted with silver nitrate in acidic media. The formula of the salt is:-

1. CaBr_2
2. $\text{Fe}_2(\text{SO}_4)_3$
3. NH_4HCO_3
4. FeBr_3
5. NH_4Br

Ans: 5

Q18) The iron (III) ion can be detected by:-

Ans: 2

1. Adding BaCl_2 solution, in acidic media and a white ppt. will appear.
2. Adding KSCN solution, and a red color will appear.
3. Adding HCl solution, a gas will change the wet red litmus paper to blue.
4. Adding NaOH solution, a gas will change the wet blue litmus paper to red.
5. Adding KSCN solution, and a white color will appear.