

دفتر إحصاء واحتمالات (كامل المادة) إعداد: ليان سليمان







denoted by t, is aux aux volume and e = 2.71828.

The probability distribution of the Poisson random variable X, representing the number of outcomes occurring in a given time interval or specified region $p(x; \lambda t) = \frac{e^{-\lambda t} (\lambda t)^x}{x!}, \quad x = 0, 1, 2, \dots,$

where λ is the average number of outcomes per unit time, distance, area, or

Table A.2 contains Poisson probability sums

$$P(r; \lambda t) = \sum_{x=0}^{r} p(x; \lambda t),$$

Talk1

for selected values of λt ranging from 0.1 to 18.0. We illustrate the use of this table Theorem 5.4: Both the mean and the variance of the Poisson distribution $p(r;\lambda t)$ are λt . During a laboratory experiment, the area distribution $p(r;\lambda t)$ are λt .

Example 5.17: During a laboratory experiment, the average number of radioactive particles pass-4 Le 1 ms 13 (s ing through a counter in 1 millisecond is 4. What is the probability that 6 particles enter the counter in a given millisecond? non bevofoutcome lintercally

Solution: Using the Poisson distribution with x = 6 and $\lambda t = 4$ and referring to Table A.2.

expected =
$$M \rightarrow \lambda t \rightarrow avgnumber = np$$

Value $f = \chi$

12 alk1

25

Example 5.19: In a certain industrial facility, accidents occur infrequently. It is known that the probability of an accident on any given day is 0.005 and accidents are independent of each other.

Pussion = Binomian (a) What is the probability that in any given period of 400 days there will be an accident on one day? \(\gamma = 1 \)

(b) What is the probability that there are at most three days with an accident?

Solution: Let X be a binomial random variable with n = 400 and p = 0.005. Thus, np = 2

Using the Poisson approximation, fcom fable(a) $P(X = 1) = e^{-2}2^1 = 0.271$ and

(b) $P(X \le 3) = \sum_{x=0}^{3} e^{-2}2^x/x! = 0.857$. 0.135335+0.270671 +0.270671 +0.180447

Example 5.20: In a manufacturing process where glass products are made, defects or bubbles

occur, occasionally rendering the piece undesirable for marketing. It is known that, on average, 1 in every 1000 of these items produced has one or more bubbles. What is the probability that a random sample of 8000 will yield fewer than 7 items In possessing bubbles?

Solution: This is essentially a binomial experiment with n = 8000 and p = 0.001. Since p is very close to 0 and n is quite large, we shall approximate with the Poisson distribution using

$$M = n\rho = \mu = (8000)(0.001) = 8.$$

Hence, if X represents the number of bubbles, we have

 $\frac{1}{2}$ $\frac{1}$

number of xample 1: A secretary makes

of Interval is the probability that on the ser page ,

what is the probability that on the ser page he or she will make x = x + 2x = 2(a) 4 or more errors? $P(x \ge 4)$ ***

(b) no errors? $P(x \ge 4)$ ***

(c) no errors? $P(x \ge 4)$ ***

(d) P(x = 0) **

(e) no errors? $P(x \ge 4)$ **

(f) P(x = 0) **

(g) P(x =


P(Y>4)=1-P(Y<4)=1-B(Y<3)

a) 0.143

P(x < 7) = P(x = 6) P(x < 7) = R(x = 7)P(x < 7) = 1 - P(x < 7)

AN Binomin distribution! b (xin, p) = (n) px q-x (q=1-p) 5000 M = np Si = npg مثلام أن دحلناها توافي Hypergeometric distribution:) (M+20) h (x3 Nonsk) = (1/2) (N-K) $\mathcal{U} = \frac{nk}{N}$ $\sigma^2 = \frac{N-n}{N-1} \cdot n \cdot \frac{K}{N} \left(1 - \frac{K}{N}\right)$ Negative Binomial distributions وجود حدادني للخاج 404057 b*(x3k,p) = (x-1) pk qx-k Geometric distributions first success g(x jp) = pgx-1 $|\mathcal{H} = \frac{1}{\rho}|_{\sigma^2} = \frac{1-\rho}{\rho^2}$ Poisson distributions

P(xint) = e (nt) 2t = 4 = np M= 02 = 2t The all are



Chapter 6

Some Continuous Probability Distributions

1




Figure 6.1: The density function for a random variable on the interval [1, 3].

$$f(1/3) = \begin{cases} \frac{1}{2} & 1 \leq 2 \leq 3 \\ 0 & 1 \leq 2 \leq 3 \end{cases}$$

- Example 6.1: Suppose that a large conference room at a certain company can be reserved for a more than 4 hours. Both long and short conferences occur quite often. In fact, it can be assumed that the length X of a conference has a uniform distribution on the interval [0, 4].
 - (a) What is the probability density function?
 - (b) What is the probability that any given conference lasts at least 3 hours?
 - Solution: (a) The appropriate density function for the uniformly distributed random variable X in this situation is

$$\begin{array}{c}
B - A \\
f(x) = \begin{cases} \frac{1}{4}, & 0 \le x \le 4, \\
0, & \text{elsewhere.} \end{cases}$$


(b)
$$P[X \ge 3] = \int_3^4 \frac{1}{4} dx = \frac{1}{4}$$
.

Theorem 6.1: The mean and variance of the uniform distribution are

$$\mu = \frac{A+B}{2} \text{ and } \sigma^2 = \frac{(B-A)^2}{12}.$$

3

Normal Distribution



Normal The density of the normal random variable X, with mean μ and variance σ^2 , is

$$n(x; \mu, \sigma) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\mathfrak{a}^2}(x-\mu)^2}, \quad -\infty < x < \infty,$$

where $\pi = 3.14159...$ and e = 2.71828....

Probability) abi other table light table I dami




Figure 6.3: Normal curves with $\underline{\underline{\mu}} < \underline{\mu}_2$ and $\underline{\sigma}_1 = \underline{\sigma}_2$.

Two normal curves having the same standard deviation but different means. The two curves are identical in form but are centered at different positions along the horizontal axis

Solving Center 1130251

5




Figure 6.4: Normal curves with $\mu_1 = \mu_2$ and $\sigma_1 < \sigma_2$.

Two normal curves with the same mean but different standard deviations

3)




Figure 6.5: Normal curves with $\mu_1 < \mu_2$ and $\sigma_1 < \sigma_2$.

symetica (1 avanute the center X-M Esse Usi (2 M-5 M+6

Two normal curves having different means and different standard deviations

Properties of the normal curve:

- The mode, which is the point on the horizontal axis where the curve is a maximum, occurs at $x = \mu$...
- The curve is symmetric about a vertical axis through the mean $\mu.$
- The curve has its points of inflection at $x = \mu \pm \sigma$; it is concave downward if μ - σ <X< μ + σ and is concave upward otherwise.
- The normal curve approaches the horizontal axis asymptotically as we proceed in either direction away from the mean.
- The total area under the curve and above the horizontal axis is 5. equal to 1.

7

Theorem 6.2: The mean and variance of $n(x; \mu, \sigma)$ are μ and σ^2 , respectively. Hence, the standard deviation is σ .

Z od X Z

Areas under the Normal Curve

The curve of any continuous probability distribution or density function is constructed so that the area under the curve bounded by the two ordinates $x = x_1$ and $x = x_2$ equals the probability that the random variable X assumes a value between $x = x_1$ and $x = x_2$. Thus, for the normal curve in Figure 6.6,

$$P(x_1 < X < x_2) = \int_{x_1}^{x_2} n(x; \mu, \sigma) \ dx = \frac{1}{\sqrt{2\pi}\sigma} \int_{x_1}^{x_2} e^{-\frac{1}{2\sigma^2}(x - \mu)^2} dx$$

is represented by the area of the shaded region.




Figure 6.6: $P(x_1 < X < x_2) = \text{area of the shaded region.}$




Figure 6.7: $P(x_1 < X < x_2)$ for different normal curve-

of a normal random variable Z with mean 0 and variance 1. This can be done by means of the transformation

$$Z = \frac{X - \mu}{\sigma}.$$

Definition 6.1: The distribution of a normal random variable with mean 0 and variance 1 is called a standard normal distribution.




Figure 6.8: The original and transformed normal distributions.

Example 6.2: Given a standard normal distribution, find the area under the curve that lies

(a) to the right of Z = 1.84 and
(b) between z = -1.97 and z = 0.86. P(Z > 1.84) = 1 - P(Z < 1.84) O = 1.84 O = 1.8

Figure 6.9: Areas for Example 6.2.

(b)

Solution: See Figure 6.9 for the specific areas.

(a)


- (a) The area in Figure 6.9(a) to the right of z = 1.84 is equal to 1 minus the area in Table A.3 to the left of z = 1.84, namely, 1 - 0.9671 = 0.0329.
- (b) The area in Figure 6.9(b) between z=-1.97 and z=0.86 is equal to the area to the left of z=0.86 minus the area to the left of z=-1.97. From Table A.3 we find the desired area to be 0.8051-0.0244=0.7807.


9

Example 6.3: Given a standard normal distribution, find the value of k such that

(a) P(Z > k) = 0.3015 and

(b) P(k < Z < -0.18) = 0.4197.





Ш

Figure 6.10: Areas for Example 6.3.

Solution: Distributions and the desired areas are shown in Figure 6.10.

- (a) In Figure 6.10(a), we see that the k value leaving an area of 0.3015 to the right must then leave an area of 0.6985 to the left. From Table A.3 it follows that(k = 0.52)
- (b) From Table A.3 we note that the total area to the left of −0.18 is equal to 0.4286. In Figure 6.10(b), we see that the area between k and -0.18 is 0.4197, so the area to the left of k must be 0.4286 - 0.4197 = 0.0089. Hence, from

Table A.3, we have k = -2.37.

A.3 table 116 to 1 has be 0.123 on 2789

Represented to the lift of 1 has be 0.123 on 2789

Represented to the lift of 1 has be 0.123 on 2789

11

Example 6.4: Given a random variable X having a normal distribution with $\mu = 50$ and $\sigma = 10$, find the probability that X assumes a value between 45 and 62.

0.5 0 Figure 6.11: Area for Example 6.4.

Solution: The z values corresponding to $x_1 = \underline{45}$ and $x_2 = 62$ are $z_1 = \frac{45 - 50}{10} = -0.5$ and $z_2 = \frac{62 - 50}{10} = 1.2$.

$$z_1 = \frac{45 - 50}{10} = -0.5$$
 and $z_2 = \frac{62 - 50}{10} = 1.2$

Therefore,

P(45 < X < 62) = P(-0.5 < Z < 1.2). P(-0.5 < Z < 1.2) is shown by the area of the shaded region in Figure 6.11. This area may be found by subtracting the area to the left of the ordinate z = -0.5from the entire area to the left of z = 1.2. Using Table A.3, we have

$$P(45 < X < 62) = P(-0.5 < Z < 1.2) = P(Z < 1.2) - P(Z < -0.5)$$

= 0.8849 - 0.3085 = 0.5764.

Given that X has a normal distribution with $\mu=300$ and $\sigma=50$, find the probability that X assumes a value greater than 362. The normal probability distribution with the desired area shaded is shown in

Figure 6.12. To find P(X > 362), we need to evaluate the area under the normal curve to the right of x = 362. This can be done by transforming x = 362 to the corresponding z value, obtaining the area to the left of z from Table A.3, and then subtracting this area from 1. We find that

$$z = \frac{362 - 300}{50} = 1.24.$$

$$P(X > 362) = P(Z > 1.24) = 1 - P(Z < 1.24) = 1 - 0.8925 = 0.1075.$$

discrete + X




Figure 6.12: Area for Example 6.5.

13

Example 6.6: Given a normal distribution with $\mu = 40$ and $\sigma = 6$, find the value of x that has

- (a) 45% of the area to the left and
- (b) 14% of the area to the right




Figure 6.13: Areas for Example 6.6.

2= -0.13

Solution: (a) An area of 0.45 to the left of the desired x value is shaded in Figure 6.13(a). We require a z value that leaves an area of 0.45 to the left. From Table A.3 we find P(Z < -0.13) = 0.45, so the desired z value is -0.13. Hence,

$$x = (6)(-0.13) + 40 = 39.22.$$

(b) In Figure 6.13(b), we shade an area equal to 0.14 to the right of the desired x value. This time we require a z value that leaves 0.14 of the area to the right and hence an area of 0.86 to the left. Again, from Table A.3, we find P(Z<1.08)=0.86, so the desired z value is 1.08 and

$$x = (6)(1.08) + 40 = 46.48.$$

0.14 auxil (1

desired lass Z auxil 9

to the we find a) ly (2

- 0.14 (2

- 0.14 (2)

- 0.14 (2)

14

Example 6.7. A certain type of storage hattery lasts, on average, 3.0 years with a deviation of 0.5 year. Assuming that battery life is normally distributed probability that a given battery will last less than 2.3 years. Solution: First construct a diagram such as Figure 6.14, showing the given distribute battery lives and the desired area. To find P(X < 2.3), we need to evaluate a area under the normal curve to the left of 2.3. This is accomplished by finding the area to the left of the corresponding z value. Hence, we find that

$$\mathcal{U}=3$$
 $z=\frac{2.3-3}{0.5}=-1.4$.
 $\mathcal{X}=2.3$ and then, using Table A.3, we have

$$P(X < 2.3) = P(Z < -1.4) = 0.0808.$$




Figure 6.14: Area for Example 6.7.

Example 6.8: An electrical firm manufactures light bulbs that have a life, before burn-out, that is normally distributed with mean equal to 800 hours and a standard deviation of 40 hours. Find the probability that a bulb burns between 778 and 834 hours.

Solution: The distribution of light bulb life is illustrated in Figure 6.15. The z values corresponding to $x_1 = 778$ and $x_2 = 834$ are

15

$$z_1 = \frac{778 - 800}{40} = -0.55$$
 and $z_2 = \frac{834 - 800}{40} = 0.85$.

P(778 < X < 834) = (P(-0.55 < Z < 0.85)) = P(Z < 0.85) - P(Z < -0.55)= 0.8023 - 0.2912 = 0.5111.




Figure 6.15: Area for Example 6.8

Example 6.10: Gauges are used to reject all components for which a certain dimension is not within the specification 1.50 \pm d. It is known that this measurement is normally \rightarrow laber \times distributed with mean 1.50 and standard deviation 0.2. Determine the value d such that the specifications "cover" 95% of the measurements. $P(X > 1.5+d) = 0.025 \Leftrightarrow P(X < 1.5+d) = 0.975$

 $P[Z \le \frac{(1.5-d)-\mu}{2}] = 0.025$ Therefore,


 $P\left[Z \le \frac{(1.5-d)-1.5}{0.20}\right] = 0.025$

from which we obtain
$$X = (1, 96 + 0.2) + 1.5$$

 $1.96 = \frac{\chi - 2.50}{0.2} = \frac{1.50 + 1.50}{0.2}$

 $P(Z \le \frac{-d}{0.20}) = 0.025$ $\mathcal{X} = 1.5 \pm d\mathcal{Y} \quad d = (0.2)(1.96) = 0.392.$

In illustration of the specifications is shown in Figure 6.17.



1-0.95

an industrial process, the diameter of a ball bearing is an important measurement. The buyer sets specifications for the diameter to be 3.0 ± 0.01 cm. The implication is that no part falling outside these specifications will be accepted. It is known that in the process the diameter of a ball bearing has a normal distribuaccepted fauge tion with mean $\mu=3.0$ and standard deviation $\sigma=0.005$. On average, how many manufactured ball bearings will be scrapped?

The distribution of diameters is illustrated by Figure 6.16. The values correspond-(2.99-3.01 ing to the specification limits are $x_1 = 2.99$ and $x_2 = 3.01$. The corresponding z $\frac{2.99 - 3.0}{0.005} = -2.0$ and $s_2 = \frac{3.01 - 3.0}{0.005} = +2.0.$ 1 - P(-2.0 < Z < 2.0) = 1 - P(Z < 2.0) + P(Z < -2.0)Hence, $= P(Z > 2.0) + P(Z < -2.0) = 2 \cdot 0.0228 = 0.0456.$ P(2.99 < X < 3.01) = P(-2.0 < Z < 2.0).From Table A.3, P(Z<-2.0)=0.0228. Due to symmetry of the normal distribu-P(z=2) - P(z=-2) tion, we find that 019772-010228 P(Z < -2.0) + P(Z > 2.0) = 2(0.0228) = 0.0456.As a result, it is anticipated that, on average, 4.56% of manufactured ball bearings will be scrapped. P(ZZ2)=P(ZZ-2)=0,0228 = 010456 0.0228 Figure 6.16: Area for Example 6.9. 17

Example 6.11: A certain machine makes electrical resistors having a mean resistance of 40 ohms and a standard deviation of 2 ohms. Assuming that the resistance follows a normal distribution and can be measured to any degree of accuracy what percentage of resistors will have a resistance exceeding 13 ohms?

Solution: A percentage is found by multiplying the relative frequency by 100%. Since the relative frequency for an interval is equal to the probability of a value falling in the interval, we must find the area to the right of x = 43 in Figure 6.18. This can be done by transforming x = 43 to the corresponding z value, obtaining the area to the left of z from Table A.3, and then subtracting this area from 1. We find $z = \frac{43 - 40}{2} = 1.5.$ Therefore,

$$P(X > 43) = P(Z > 1.5) = 1 - P(Z < 1.5) = 1 - 0.9332 = 0.0668.$$

Hence, 6.68% of the resistors will have a resistance exceeding 43 ohms.





Figure 6.18: Area for Example 6.11.

ces exceeding 43 ohms for Example 6.11 if resistance Solution: This problem differs from that in Example 6.11 in that we now assign a measured for the nearest olum.

Solution: This problem differs from that in Example 6.11 in that we now assign a measure-ment of 43 ohms to all resistors whose resistances are greater than 42.5 and less ment of 43.5. We are actually approximating a discrete distribution by means of a continuous normal distribution. The required area is the region shaded to the right of 43.5 in Figure 6.19. We now find that $z = \frac{43.5 - 40}{9} = 1.75.$ Hence. P(X > 43.5) = P(Z > 1.75) = 1 - P(Z < 1.75) = 1 - 0.9599 = 0.0401.



Therefore, 4.01% of the resistances exceed 43 ohms when measured to the nearest ohm. The difference 6.68%-4.01%=2.67% between this answer and that of Example 6.11 represents all those resistance values greater than 43 and less than 43.5 that are now being recorded as 43 ohms.




Figure 6 10; Area for Example 6.12

19

Example 6.13: The average grade for an exam is 74, and the standard deviation is 7. If 12% of the class is given As, and the grades are curved to follow a normal distribution, what is the lowest possible A and the highest possible B?

Solution: In this example, we begin with a known area of probability, find the z value, and then determine x from the formula $x = \sigma z + \mu$. An area of 0.12, corresponding to the fraction of students receiving As, is shaded in Figure 6.20. We require a z value that leaves 0.12 of the area to the right and, hence, an area of 0.88 to the left. From Table A.3, P(Z < 1.18) has the closest value to 0.88, so the desired z






Figure 6.20: Area for Example 6.13.

74



74 De

0.6



and Exponential Distributions

The gamma function is defined by

$$\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$$
, for $\alpha > 0$.

The following are a few simple properties of the gamma function.

(a) $\Gamma(n) = (n-1)(n-2)\cdots(1)\Gamma(1)$, for a positive integer n.

To see the proof, integrating by parts with $u = x^{\alpha-1}$ and $dv = e^{-x} dx$, we obtain

$$\Gamma(\alpha) = -e^{-x} |x^{\alpha - 1}|_0^{\infty} + \int_0^{\infty} e^{-x} (\alpha - 1) x^{\alpha - 2} |dx| = (\alpha - 1) \int_0^{\infty} x^{\alpha - 2} e^{-x} |dx|.$$

for $\alpha > 1$, which yields the recursion formula

$$\Gamma(\alpha) = (\alpha - 1)\Gamma(\alpha - 1).$$

The continuous $\$ random variable X has a gamma distribution, with parameters α and β , if its density function is given by

where $\alpha > 0$ and $\beta > 0$. $\begin{cases}
f(x; \alpha, \beta) = \begin{cases} \frac{1}{\beta^{\alpha} \Gamma(\alpha)} x^{\alpha - 1} e^{-x/\beta}, & x > 0, \\ 0, & \text{elsewhere,} \end{cases}$ $S(\alpha) \notin \text{Parameter} \quad \text{Where } (\alpha) = (n - 1)!$ $(\alpha \lor \beta) \quad (\alpha \lor$

s exporential

Function specific number of events is the parameter α in the gamma density function

a (alpha) is known as the shape parameter, while b (beta) is referred to as the scale parameter





Figure 6.28: Gamma distributions.

fime) ((() () () () ()

The continuous random variable X has an exponential distribution, with Distribution parameter β , if its density function is given by

 $f(x;\beta) = \begin{cases} \frac{1}{\beta}e^{-x/\beta}, & x > 0, \\ 0, & \text{elsewhere.} \end{cases}$ where $\beta > 0$.

1.0



modelling for time -> gamma


gamma de Ricer lo

Suppose that telephone calls arriving at a particular switchboard follow a Poisson process with an average of 5 calls coming per minute. What is the probability that up to a minute will clapse by the time 2 calls have come in to the switchboard? Solution: The Poisson process applies, with time until 2 Poisson events following a gamma distribution with $\beta = 1/5$ and $\alpha = 2$. Denote by X the time in minutes that transpires before 2 calls come. The required probability is given by $P(X \le 1) = \binom{1}{2}$

While the origin of the gamma distribution deals in time (or space) until the occurrence of α Poisson events, there are many instances where a gamma distribution works very well even though there is no clear Poisson structure. This is particularly true for survival time problems in both engineering and biomedical applications.

the mean of the exponential distribution is the parameter 0, the reciprocal of the parameter in the Poisson distribution

B = Toisson



chample 6.21: Consider Exercise 3.31 on page 94. Based on extensive testing, it is determined that the time Y in years before a major repair is required for a certain washing machine is characterized by the density function

$$f(y) = \begin{cases} \frac{1}{4}e^{-y/4}, & y \ge 0, \\ 0, & \text{elsewhere.} \end{cases}$$

Solution: Consider the cumulative distribution function F(y) for the exponential distribution,

$$F(y) = \frac{1}{\beta} \int_0^y e^{-t/\beta} dt = 1 - e^{-y/\beta},$$

Then


$$P(Y > 6) = 1 - F(6) = e^{-3/2} = 0.2231.$$

Thus, the probability that the washing machine will require major repair after year six is 0.223. Of course, it will require repair before year six with probability 0.777. Thus, one might conclude the machine is not really a bargain. The probability that a major repair is necessary in the first year is

$$P(Y < 1) = 1 - e^{-1/4} = 1 - 0.779 = 0.221.$$

п

25



3.31 Based on extensive testing, it is determined by the manufacturer of a washing machine that the time Y (in years) before a major repair is required is characterized by the probability density function

$$f(x) = \begin{cases} \frac{1}{1}e^{-\frac{y}{4}} & y \ge 0, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Critics would certainly consider the product a bargain if it is unlikely to require a major repair before the sixth year. Comment on this by determining P(Y > 6).
- (b) What is the probability that a major repair occurs in the first year?
- (a) For $y \ge 0$, $F(y) = \frac{1}{4} \int_0^y e^{-t/4} dy = 1 e^{y/4}$. So, $P(Y > 6) = e^{-6/4} = 0.2231$. This probability certainly cannot be considered as "unlikely."
- (b) $P(Y \le 1) = 1 e^{-1/4} = 0.2212$, which is not so small either.

20

The continuous random variable X has a chi-squared distribution, with vDistribution degrees of freedom, if its density function is given by

 $f(x;v) = \begin{cases} \frac{1}{2^{c/2}\Gamma(v/2)} x^{v/2-1} e^{-x/2}, & x > 0, \\ 0, & \text{elsewhere,} \end{cases}$

Theorem 6.5: The mean and variance of the chi-squared distribution are

27

Definition 6.3: A beta function is defined by

$$B(\alpha,\beta) = \int_0^1 x^{\alpha-1} (1-x)^{\beta-1} dx = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, \text{ for } \alpha,\beta > 0,$$

where $\Gamma(\alpha)$ is the gamma function.

Bera Distribution The continuous random variable X has a beta distribution with parameters $\alpha > 0$ and $\beta > 0$ if its density function is given by

$$f(x) = \begin{cases} \frac{1}{B(\alpha,\beta)} x^{\alpha-1} (1-x)^{\beta-1}, & 0 < x < 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Note that the uniform distribution on (0,1) is a beta distribution with parameters (n-1)! = 7! $\alpha = 1$ and $\beta = 1$.

The mean and variance of a beta distribution with parameters α and β are

Theorem 6.6:

$$\mu = \frac{\alpha}{\alpha + \beta}$$
 and $\sigma^2 = \frac{\alpha \beta}{(\alpha + \beta)^2 (\alpha + \beta + 1)}$,

respectively.

For the uniform distribution on (0,1), the mean and variance are

$$\mu = \frac{1}{1+1} = \frac{1}{2} \text{ and } \sigma^2 = \frac{(1)(1)}{(1+1)^2(1+1+1)} = \frac{1}{12},$$

$$\mathcal{U} \quad \text{w.}$$

The continuous random variable X has a lognormal distribution if the random variable $Y = \ln(X)$ has a normal distribution with mean μ and standard deviation σ . The resulting density function of X is

$$f(x;\mu,\sigma) = \begin{cases} \frac{1}{\sqrt{2\pi}\sigma x} e^{-\frac{1}{2\sigma^2}|\ln(x)-\mu|^2}, & x \ge 0, \\ 0, & x < 0. \end{cases}$$




Figure 6.29: Lognormal distributions

The mean and variance of the lognormal distribution are

$$\mu = e^{\mu + \sigma^2/2}$$
 and $\sigma^2 = e^{2\mu + \sigma^2} (e^{\sigma^2} - 1)$.

In clog) (8his) normal distribution

29

Example 6.22: Concentrations of pollutants produced by chemical plants historically are known to exhibit behavior that resembles a lognormal distribution. This is important when one considers issues regarding compliance with government regulations. Suppose it is assumed that the concentration of a certain pollutant, in parts per million, has a lognormal distribution with parameters $(\mu = 3.2)$ and $(\sigma = 1)$ What is the probability that the concentration exceeds 8 parts per million? الجادزاري

Solution: Let the random variable X be pollutant concentration. Then

$$P(X > 8) = 1 - P(X \le 8)$$

Since ln(X) has a normal distribution with mean $\mu = 3.2$ and standard deviation

$$P(X \le 8) = \Phi \left[\frac{\ln(8) - 3.2}{1} \right] = \Phi(-1.12) = 0.1314. \quad -0.1314$$

$$= 0.8686$$

example 6.23: The life, in thousands of miles, of a certain type of electronic control for locomotives has an approximately lognormal distribution with $\mu = 5.149$ and $\sigma = 0.737$. Find

Solution: From Table A.3, we know that P(Z < -1.645) = 0.05. Denote by X the life of such an electronic control. Since $\ln(X)$ has a part of the life of such an electronic control. $\mu = 5.149$ and $\sigma = 0.737$, the 5th percentile of X can be calculated as

$$\ln(x) = 5.149 + (0.737)(-1.645) = 3.937.$$

Hence, x = 51.265. This means that only 5% of the controls will have lifetimes less. than 51,265 miles.

The continuous random variable X has a Weibull distribution, ψ Distribution eters α and β , if its density function is given by

and
$$\beta$$
, if its density function is given by
$$f(x; \alpha, \beta) = \begin{cases} \alpha \beta x^{\beta - 1} e^{-\alpha x^{\beta}}, & x > 0, \\ 0, & \text{elsewhere,} \end{cases}$$

where $\alpha > 0$ and $\beta > 0$.

Theorem 6.8: The mean and variance of the Weibull distribution are

The mean and variance of the vicinia
$$\mu = \alpha^{-1/\beta} \Gamma\left(1 + \frac{1}{\beta}\right) \text{ and } \sigma^2 = \alpha^{-2/\beta} \left\{ \Gamma\left(1 + \frac{2}{\beta}\right) - \left[\Gamma\left(1 + \frac{1}{\beta}\right)\right]^2 \right\}.$$

31




Figure 6.30: Weibull distributions ($\alpha = 1$).

The cumulative distribution function for the Weibull distribution is given by

$$F(x) = 1 - e^{-\alpha x^{\beta}}, \quad \text{for } x \ge 0,$$
 for $\alpha > 0$ and $\beta > 0$.

Example 6.24: The length of life X, in hours, of an item in a machine shop has a Weibull distribution with $\alpha = 0.01$ and $\beta = 2$. What is the probability that it fails before eight / hours of usage?

Solution:
$$P(X < 8) = F(8) = 1 - e^{-(0.01)8^2} = 1 - 0.527 = 0.473$$
.

In a certain city, the daily consumption of electric power, in millions of kilowatt-hours, is a random variable X having a gamma distribution with $\mu = 6$ and variance $\sigma^2 = 12$.

- (a) Find the values of α and β .
- (b) Find the probability that on any given day the daily power consumption will exceed 12 million kilowatthours. $\rho(x>12)$

33

$$\beta = \left(\frac{6}{\alpha}\right)$$

[XB=6]-0

 $\alpha = \frac{36}{36} = 3$ $\alpha = \frac{36}{3} = 3$ $\alpha = \frac{36}{3} = 3$

· Answer:

(a)
$$\mu = \alpha\beta = 6$$

$$\sigma^2 = \alpha \beta^2 = 12$$

So, $\beta = 2$ and then $\alpha = 3$.

(b)
$$P(X > 12) = \frac{1}{16} \int_{12}^{\infty} x^2 e^{-x/2} dx$$

$$P(X > 12) = \frac{1}{16} \left[-2x^2 e^{-x/2} - 8xe^{-x/2} - 16e^{-x/2} \right]_{12}^{\infty} = 25e^{-6} = 0.0620$$

11,000

· The length of time for one individual to be served at a cafeteria is a random variable having an exponential distribution with a mean of 4 minutes. What is the probability that a person is served in less than 3 minutes on at least 4 of the next 6 days?) Birowal

B=4

$$P(X < 3) = \frac{1}{4} \int_0^3 e^{-x/4} dx = -e^{-x/4} \Big|_0^3 = 1 - e^{-3/4} = 0.5276.$$

$$P(Y \ge 4) = \sum_{x=4}^{6} b(y; 6, 1 - e^{-3/4}) = {6 \choose 4} (0.5276)^4 (0.4724)^2 + {6 \choose 5} (0.5276)^5 (0.4724) + {6 \choose 6} (0.5276)^6 = 0.3968.$$

$$P(x \le 3) = \int \frac{1 - x/4}{4} dx = -\frac{x}{4} e^{-\frac{x}{4}} = -\frac{x}{4} e^{-\frac{x}{4}}$$

$$= 0.5276$$

$$P(x \ge 4) = 1 - P(x \ge 4)$$

$$= 1 - 0.0563 = 0.3437$$

$$1.1553$$

End of Chapter 6


1) finally

Chi form distribution

$$F(x) \land I(B) = \begin{cases} 1 \\ B \land A \end{cases}$$

$$A \in X \neq B$$





Probability and Statistics

Chapter 5

1

Chapter 5


Some Discrete Probability
Distributions

Discrete Uniform Distribution

Example 5.1: When a light bulb is selected at random from a box that contains a 40-watt bulb a 60-watt bulb, a 75-watt bulb, and a 100-watt bulb, each element of the sample space S = {40, 60, 75, 100} occurs with probability 1/4. Therefore, we have a uniform distribution, with

> المناور ا ما المرابع المر of sumple space foind

Example 5.2: When a fair die is tossed, each element of the sample space $S = \{1,2,3,4,5,6\}$ occurs with probability 1/6. Therefore, we have a uniform distribution, with






Figure 5.1: Histogram for the tossing of a die

The Bernoulli Process)

Strictly speaking, the Bernoulli process must possess the following properties:

The experiment consists of repeated trials.

2. Each trial results in an outcome that may be classified as a success or a failure.

3. The probability of success, denoted by p, remains constant from trial to trial.

The repeated trials are independent.


Consider the set of Bernoulli trials where three items are selected at random from a manufacturing process, inspected, and classified as defective or nondefective. A defective item is designated a success. The number of successes is a random variable X assuming integral values from 0 through 3. The eight possible outcomes and the corresponding values of X are

Outcome NDN | NND Since the items are selected independently and we assume that the process produces 25% defectives, we have

 $P(NDN) = P(N)P(D)P(N) = \begin{pmatrix} \frac{3}{4} \end{pmatrix} \begin{pmatrix} \frac{1}{4} \end{pmatrix} \begin{pmatrix} \frac{3}{4} \end{pmatrix} = \frac{9}{64}.$ $P(NDN) = P(N)^{\frac{3}{4}} - \frac{3}{4} \begin{pmatrix} \frac{3}{4} \end{pmatrix} = \frac{9}{64}.$

Similar calculations yield the probabilities for the other possible outcomes. The probability distribution of X is therefore

حالمِل جمع حالمِل جمع كل اللي عدم واحر لكي عنزهم و



Outcome	Probability	X			
NNN.	$\frac{3}{4} \times \frac{3}{4} \times \frac{3}{4} = \frac{27}{64}$	0			
NND	3 3 1 9	1	The prob	The probability distribution of X is	
	4 4 4 64	-	Z.	f(x)=P(X=	
NDN	3 1 3 9	1	0	27 64	
I A : A	4 4 4 64		pu 1	$\frac{9}{64} + \frac{9}{64} + \frac{9}{64}$	
NDD	$\frac{3}{-} \times \frac{1}{-} \times \frac{1}{-} = \frac{3}{-}$	2	2	3 3 3	
	4 4 4 64			64 64 64	
DNN	$\frac{1}{4} \times \frac{3}{4} \times \frac{3}{4} = \frac{9}{64}$	1	3	64	
	4 4 4 64				
DND	$\frac{1}{-\times}\frac{3}{\times}\frac{1}{\times}=\frac{3}{-}$	2			
	4 4 4 64				
DDN	$\frac{1}{4} \times \frac{1}{4} \times \frac{3}{4} = \frac{3}{4}$	2			
	4 4 4 64				
DDD	$\frac{1}{\times}$ $\frac{1}{\times}$ $\frac{1}{\times}$ $\frac{1}{\times}$ $\frac{1}{\times}$	3			
31 34 3	4 4 4 64				


The number X of successes in n Bernoulli trials is called a binomial random variable. The probability distribution of this discrete random variable is called the binomial distribution, and its values will be denoted by b(x; n, p) since they depend on the number of trials and the probability of a success on a given trial, Thus, for the probability distribution of X, the number of defectives is

$$P(X = 2) = f(2) = b\left(2; 3, \frac{1}{4}\right) = \frac{9}{64}.$$

A Bernoulli trial can result in a success with probability p and a failure with probability q = 1 - p. Then the probability distribution of the binomial random variable X, the number of successes in n independent trials, is

Note that when
$$n=3$$
 and $p=1/4$, the probability distribution of X , the number of defectives, may be written as

$$b\left(x;3,\frac{1}{4}\right) = \binom{3}{x}\left(\frac{1}{4}\right)^x\left(\frac{3}{4}\right)^{3-x}, \quad x = 0,1,2,3,$$
rather than in the tabular form on page 144.



п

Example 5.1: The probability that a certain kind of component will survive a shock test if 3/4.] Find the probability that exactly 2 of the next 4 components tested survive. Solution: Assuming that the tests are independent and p=3/4 for each of the 4 tests, we

$$\rho = \frac{55\sqrt{\frac{3}{4}}}{\frac{3}{4}} \qquad b\left(2;4,\frac{3}{4}\right) = \left(\frac{4}{2}\right)\left(\frac{3}{4}\right)^{2}\left(\frac{1}{4}\right)^{2} = \left(\frac{4!}{2! \ 2!}\right)\left(\frac{3^{2}}{4^{4}}\right) = \frac{27}{128}.$$

are known to have contracted this disease, what is the probability that (a) at least 10 survive, (b) from 3 to 8 survive, and (c) exactly 5 survive? Solution: Let X be the number of people who survive (a) $P(X \ge 10) = 1 - P(X < 10) = 1 - \sum_{x \ge 10} b(x; 15, 0.4) = 1 - 0.9662$ (b) $P(3 \le X \le 8) = \sum b(x; 15, 0.4) = \sum b(x; 15, 0.4) - \sum b(x; 15, 0.4)$ = 0.9050 - 0.0271 = 0.8779 $P(X = 5) = b(5; 15, 0.4) = \sum_{x=0}^{\infty} b(x; 15, 0.4) - \sum_{x=0}^{\infty} b(x; 15, 0.4)$) f(5) - f(4) =0.1859

The probability that a patient recovers from a rare blood discuse is 0.4. If 15 people

no 15

1210

The probability that a patient recovers from a rare blood disease is 0.4. If 15 people are known to have contracted this disease, what is the probability that (a) at least 10 survive, (b) from 3 to 8 survive, and (c) exactly 5 survive?

Solution: Let X be the number of people who survive. $b(x; n, p) = \binom{n}{x} p^x q^{n-x}$

 $P(X \ge 10)$ P(X=10)+P(X=11)+P(X=12)+P(X=13)+P(X=14)+P(X=15)

 $\binom{18}{10} (0.4)^{10} (0.6)^{2} + \binom{16}{11} (0.4)^{11} (0.6)^{4} + \binom{16}{12} (0.4)^{12} (0.6)^{2} + \binom{16}{11} (0.4)^{13} (0.6)^{2} + \binom{16}{14} (0.6)^{2} + \binom{18}{14} (0.6)^{2}$

(b) $P(3 \le X \le 8) = P(X=3)+P(X=4)+P(X=5)+P(X=6)+P(X=7)+P(X=8)$

 $\binom{15}{3} (0.4)^3 (0.6)^{12} + \binom{15}{4} (0.4)^4 (0.6)^{11} + \binom{15}{5} (0.4)^5 (0.6)^{11} + \binom{15}{6} (0.4)^6 (0.6)^{11} + \binom{15}{7} (0.4)^7 (0.6)^8 + \binom{15}{8} (0.4)^8 (0.6)^7$

(C) (P X=5)= $\binom{15}{5}$ (0.4)⁵ (0.6)¹⁰ = 0.1859

1) P(XZIO) = 1- P(XZIO) - 1-0,9652

inple 5.3: A large chain retailer purchases a certain kind of electronic device from a manufacturer. The manufacturer indicates that the defective rate of the device is 3%.

(a) The inspector randomly picks 20 items from a shipment. What is the proba-((X21) bility that there will be at least one defective item among these 20?

(b) Suppose that the retailer receives 10 shipments in a month and the inspector randomly tests 20 devices per shipment. What is the probability that there will be exactly 3 shipments each containing at least one defective device among the 20 that are related by the containing at least one defective device among the 20 that are selected and tested from the shipment?

Solution: (a) Denote by X the number of defective devices among the 20. Then X follows

(a) Denote by
$$X$$
 the Hamber of detection. Hence, $\rho(X \ge 1)$

$$P(X \ge 1) = 1 - P(X = 0) = 1 - b(0; 20, 0.03)$$

$$= 1 - (0.03)^{0}(1 - 0.03)^{20-0} = 0.4562.$$

(b) In this case, each shipment can either contain at least one defective item or not. Hence, testing of each shipment can be viewed as a Bernoulli trial with p = 0.4562 from part (a). Assuming independence from shipment to shipment and denoting by Y the number of shipments containing at least one defective item, Y follows another binomial distribution b(y; 10, 0.4562). Therefore,

Example 5.4: It is conjectured that an impurity exists in 30% of all drinking wells in a certain rural community. In order to gain some insight into the true extent of the problem, it is determined that some testing is necessary. It is too expensive to test all of the wells in the area, so 10 are randomly selected for testing.

(a) Using the binomial distribution, what is the probability that exactly 3 wells have the impurity, assuming that the conjecture is correct?

(b) What is the probability that more than 3 wells are impure?

Solution: (a) We require $b(3; 10, 0.3) = \sum_{x=0}^{3} b(x; 10, 0.3) - \sum_{x=0}^{2} b(x; 10, 0.3) = 0.6496 - 0.3828 = 0.2668.$ $h = \begin{cases} 0 & \text{if } x = 0 \\ 0 & \text{if } x = 0 \end{cases}$ (b) In this case, P(X > 3) = 1 - 0.6496 = 0.3504.(a) $\binom{10}{3}$ $(0.3)^3$ $(0.7)^7 = 0.2668$ $P(X > 3) = 1 - P(X \le 3)$ = 1 - 0, 6.96

1-{0.0282+0.1211+0.2335+0.2668}

Since the probability distribution of any binomial random variable depends only on the values assumed by the parameters n, p, and q, it would seem reasonable to assume that the mean and variance of a binomial random variable also depend on the values assumed by these parameters. Indeed, this is true, and in the proof of Theorem 5.1 we derive general formulas that can be used to compute the mean and variance of any binomial random variable as functions of n, p, and q.

Theorem 5.1:

The mean and variance of the binomial distribution b(x; n, p) are $\mu = np \text{ and } \sigma^2 = npq.$

13

Example 5.5: Find the mean and variance of the binomial random variable of Example 5.2, and then use Chebyshev's theorem (on page 137) to interpret the interval $\mu \pm 2\sigma$.

Solution: Since Example 5.2 was a binomial experiment with n = 15 and p = 0.4, by Theorem

5.1, we have $\eta \quad \rho \quad \eta \quad \eta \quad \rho \quad (1 - P)$ $\mu = (15)(0.4) = 6 \text{ and } \sigma^2 = (15)(0.4)(0.6) = 3.6.$

Taking the square root of 3.6, we find that $\sigma = 1.897$. Hence, the required interval is $6\pm(2)(1.897)$, or from 2.206 to 9.794. Chebyshev's theorem states that the number of recoveries among 15 patients who contracted the disease has a probability of at least 3/4 of falling between 2.206 and 9.794 or, because the data are discrete, between 2 and 10 inclusive.

There are solutions in which the computation of binomial probabilities may allow us to draw a scientific inference about population after data are collected. An illustration is given in the next example.

are known to have contracted this disease, what is the probability that (a) at least 10 shrvive (b) from 3 to 8 survive, and (c) exactly 5 survive?

Example 5.6: Consider the situation of Example 5.4. The notion that 30% of the wells are impure is merely a conjecture put forth by the area water board. Suppose 10 wells are randomly selected and 6 are found to contain the impurity. What does this imply about the conjecture? Use a probability statement.

Solution: We must first ask: "If the conjecture is correct, is it likely that we would find 6 or more impure wells?"

$$= 1 - \rho (x < 6) \frac{10}{10}$$

$$P(X \ge 6) = \sum_{x=0}^{5} b(x; 10, 0.3) - \sum_{x=0}^{5} b(x; 10, 0.3) = 1 - 0.9527 = 0.0473.$$

As a result, it is very unlikely (4.7% chance) that 6 or more wells would be found impure if only 30% of all are impure. This casts considerable doubt on the conjec-

ture and suggests that the impurity problem is much more severe.

As the reader should realize by now, in many applications there are more than two possible outcomes. To borrow an example from the field of genetics, the color of guinea pigs produced as offspring may be red, black, or white. Often the "defective" or "not defective" dichotomy is truly an oversimplification in engineering situations. Indeed, there are often more than two categories that characterize items or parts coming off an assembly line.

15

Suppose a large urn contains 400 red marbles and 600 blue marbles.

A random sample of 10 marbles is drawn without replacement. What is the probability exactly 3 are red?

$$P(X=3) = {\binom{400}{3}} {\binom{600}{7}} = 0.2155$$

Ivpergeometric Distribution) ~ grown 5-9 500

Distribution

The probability distribution of the hypergeometric random variable X, the number of successes in a random sample of size n selected from N items of which k are labeled success and N-k labeled failure, is

$$h(x;N,n,k) = \frac{\binom{k}{x}\binom{N-k}{n-x}}{\binom{N}{n}}, \quad \max\{0,n-(N-k)\} \le x \le \min\{n,k\}.$$


Example 5.9: Lots of 40 components each are deemed unacceptable if they contain 3 or more defectives. The procedure for sampling a lot is to select 5 components at random and to reject the lot if a defective is found. What is the probability that exactly 1 defective is found in the sample if there are 3 defectives in the entire lot?

Solution: Using the hypergeometric distribution with n = 5, N = 40, k = 3, and x = 1, we find the probability of obtaining 1 defective to be

40 15 411

$$h(1;40,5,3) = \frac{\binom{3}{1}\binom{37}{4}}{\binom{40}{5}} = 0.3011.$$

Once again, this plan is not desirable since it detects a bad lot (3 defectives) only about 30% of the time.



17

Exercise 5.31

A homeowner plants 6 bulbs selected at random from a box containing 5 tulip bulbs and 4 daffodil bulbs. What is the probability that he planted 2 daffodil bulbs and 4 tulip bulbs?

$$h(2; 9, 6, 4) = \frac{\binom{4}{2}\binom{5}{4}}{\binom{9}{6}} = \frac{5}{14}.$$

Exercise 5.32

From a lot of 10, 4 are selected at random and fired. If the lot contains 3 defective missiles that will not fire, what is the probability that (a) all 4 will fire? (b) at most 2 will not fire?

(a)
$$h(4;10,4,7) = 1/6$$
(b) $\sum_{i=1/6}^{2} h(x;10,4,3) = \frac{29}{30}$

(a)
$$h(4;10,4,7) = 1/6$$

(b) $\sum_{x=0}^{2} h(x;10,4,3) = \frac{29}{30}$

$$\frac{\binom{3}{2}\binom{7}{2}}{\binom{4}{4}} = \frac{3}{10} \left(\begin{array}{c} \binom{7}{4}\binom{3}{4} = 1/6 \\ \binom{7}{4}\binom{7}{4} + \binom{3}{1}\binom{7}{3}\binom{7}{3} + \binom{3}{2}\binom{7}{4}\binom{7}{4} + \binom{3}{1}\binom{7}{3}\binom{7}{4}\binom{7}\binom{7}{4}\binom{7}{4}\binom{7}{4}\binom{7}$$

Theorem 5.2: The mean and variance of the hypergeometric distribution h(x; N, n, k) are

$$\mu = \frac{nk}{N} \text{ and } \sqrt{2} = \frac{N-n}{N-1} \cdot n \cdot \frac{k}{N} \left(1 - \frac{k}{N} \right).$$

Example 5.10: Let us now reinvestigate Example 3.4 on page 83. The purpose of this example was to illustrate the notion of a random variable and the corresponding sample space. In the example, we have a lot of 100 items of which 12 are defective. What is the probability that in a sample of 10, 3 are defective?

Solution: Using the hypergeometric probability function, we have

$$h(3;100,10,12) = \frac{\binom{12}{3}\binom{88}{7}}{\binom{100}{10}} = 0.08. \qquad \mathcal{H} = \underbrace{12 + 107}_{100} = 1.2$$

Example 3.4: Statisticians use sampling plans to either accept or reject batches or lots of material. Suppose one of these sampling plans involves sampling independently 10 items from a lot of 100 items in which 12 are defective,

Let X be the random variable defined as the number of items found defective in the sample of 10. In this case, the random variable takes on the values 0, 1, 2, ..., 9, 10,

$$\sigma^{2} = \frac{100 - 10}{100 - 1} \cdot 10 + 4 \frac{12}{100} \left(1 - \frac{12}{100}\right) = \frac{24}{25}$$

1: Find the mean and variance of the random variable of Example 5.9 and then use Chebyshev's theorem to interpret the interval $\mu \pm 2\sigma$.

Solution: Since Example 5.9 was a hypergeometric experiment with N = 40, n = 5, and k = 3, by Theorem 5.2, we have

$$\mu = \frac{(5)(3)}{40} = \frac{3}{8} = 0.375,$$

$$\sigma^2 = \left(\frac{40 - 5}{39}\right)(5)\left(\frac{3}{40}\right)\left(1 - \frac{3}{40}\right) = 0.3113.$$

Taking the square root of 0.3113, we find that $\sigma = 0.558$. Hence, the required interval is $0.375 \pm (2)(0.558)$, or from -0.741 to 1.491. Chebyshev's theorem states that the number of defectives obtained when 5 components are selected at random from a lot of 40 components of which 3 are defective has a probability of at least 3/4 of falling between -0.741 and 1.491. That is, at least three-fourths of the time, the 5 components include fewer than 2 defectives.

Example 5.9: Lots of 40 components each are deemed unacceptable if they contain 3 or more defectives. The procedure for sampling a lot is to select 5 components at random and to reject the lot if a defective is found. What is the probability that exactly 1 defective is found in the sample if there are 3 defectives in the entire lot?

Solution: Using the hypergeometric distribution with n = 5, N = 40, k = 3, and x = 1, we find the probability of obtaining 1 defective to be

21

$$h(1;40,5,3) = \frac{\binom{3}{1}\binom{37}{4}}{\binom{60}{5}} = 0.3011.$$

If repeated independent trials can result in a success with probability p and number of a failure with probability q = 1 - p, then the probability distribution of the random variable X, the number of the trial on which the kth success occurs, is kth success

occurs on the xth trial. Experiments of $(x,k,p) = {x-1 \choose k-1} p^k q^{x-k}$, $x=k,k+1,k+2,\ldots$ experiments.

Example 5.14: In an NBA (National Basketball Association) championship series, the team that wins four games out of seven is the winner. Suppose that teams A and B face each other in the championship games and that team A has probability 0.55 of winning a game over team B.

(a) What is the probability that team A will win the series in 6 games?

(b) What is the probability that team A will win the series?

(c) If teams A and B were facing each other in a regional playoff series, which is decided by winning three out of five games, what is the probability that team A would win the series?

Solution: (a) $b^*(6; 4, 0.55) = \binom{5}{4} \cdot 0.55^4 \cdot (1 - 0.55)^{6-4} = 0.1853$

(b) P(team A wins the championship series) is

$$b^*(4;4,0.55) + b^*(\underline{5;4,0.55}) + b^*(\underline{6;4,0.55}) + b^*(\underline{7;4,0.55})$$

$$= 0.0915 + 0.1647 + 0.1853 + 0.1668 = 0.6083.$$

(c) P(team A wins the playoff) is

$$b^*(3; 3, 0.55) + b^*(4; 3, 0.55) + b^*(5; 3, 0.55)$$

= $0.1664 + 0.2246 + 0.2021 = 0.5931$.

SUCTUSE

If repeated independent trials can result in a success with probability p and a failure with probability q = 1 - p, then the probability distribution of the random variable X, the number of the trial on which the first success occurs, is

الوقدا عمادة المية الخرهة بم التحاح

 $g(x;p) = pq^{x-1}, \quad x = 1, 2, 3, \dots$

ample 5.15: For a certain manufacturing process, it is known that, on the average, 1 in every 100 items is defective. What is the probability that the fifth item inspected is the first defective item found? ution: Using the geometric distribution with x = 5 and p = 0.01, we have

 $g(5;0.01) = (0.01)(0.99)^4 = 0.0096$. ztill Go delective 10.1

Example 5.16: At a "busy time," a telephone exchange is very near capacity, so callers have difficulty placing their calls. It may be of interest to know the number of attempts necessary in order to make a connection. Suppose that we let p = 0.05 be the probability of a connection during a busy time. We are interested in knowing the probability that 5 attempts are necessary for a successful call.

Solution: Using the geometric distribution with x = 5 and p = 0.05 yields $P(X = x) = g(5; 0.05) = (0.05)(0.95)^4 = 0.041.$

Quite often, in applications dealing with the geometric distribution, the mean and variance are important. For example, in Example 5.16, the expected number of calls necessary to make a connection is quite important. The following theorem states without proof the mean and variance of the geometric distribution.

23

Theorem 5.3: The mean and variance of a random variable following the geometric distribution are

$$\mu = \frac{1}{p} \text{ and } \sigma^2 = \frac{1-p}{p^2}.$$

distribution of the number of successes in a fixed multiper of independent Bernoulli

distribution of the number of trials needed to get a fixed number of successes

لمان سليمان احمد ياست



Probability and Statistics

Chapter 3

أنواع المتغيرات الهشوالية وله متعل لا منعمل

1

Chapter 3

Random Variables and Probability
Distributions

a rabulately many

3 in atilles

Three electrical components are tested:

 $S = \{NNN, NND, NDN, DNN, NDD, DND, DDN, DDD\}$

we are concerned with the number of

defectives that occur?

0,1,2, or 3 متنوعلواع : رقم معيقى E = {DDN, DND, NDD}

sample Ilpivill * SPACE

(3) Just yet 4 2= (011/2)

7-> nonky of black

Definition 3.1: A random variable is a function that associates a real number with each element in the sample space.

Example 3.1: Two balls are drawn in succession without replacement from an urn containing 4 red balls and 3 black balls. The possible outcomes and the values y of the random variable Y, where Y is the number of red balls, are

5= [KB/ KB/BR/ BB} (3) get 1 yet 1 4 9 = 0/1/2


Example 3.2: A stockroom clerk returns three safety helmets at random to three steel mill employees who had previously checked them. If Smith, Jones, and Brown in that order) receive one of the three hats, list the sample points for the possible orders of returning the helmets, and find the value m of the random variable M that represents the number of correct matches.


Solution: If S, J, and B stand for Smith's, Jones's, and Brown's helmets, respectively, then the possible arrangements in which the helmets may be returned and the number

of correct matches are


العلقة اللي المتعلقة اللي المتعلقا JSB JBSBSJ


() AL () 12 4 () 14 () 19





- Example 3.6: Interest centers around the proportion of people who respond to a certain mail order solicitation. Let X be that proportion. X is a random variable that takes on all values x for which $0 \le x \le 1$. П
- Definition 3.2: If a sample space contains a finite number of possibilities or an unending sequence with as many elements as there are whole numbers, it is called a discrete sample space.
- Definition 3.3: If a sample space contains an infinite number of possibilities equal to the number of points on a line segment, it is called a continuous sample space.





I Berobublity

side airbags among the next 4 cars sold by the agency/the number of ways of selling 3 cars with side airbags. Solution: Since the probability of selling an automobile with side airbags is 0.5, the 2⁴ = 16 points in the sample space are equally likely to occur. Therefore, the denominator for all probabilities, and also for our function, is 16. To obtain the number of ways of selling 3 cars with side airbags, we need to consider the number of ways of partitioning 4 outcomes into two cells, with 3 cars with side airbags assigned to one cell and the model without side airbags assigned to the other. This can be done in $\binom{4}{3}$ = 4 ways. In general, the event of selling x models with side airbags and 4-x models without side airbags can occur in $\binom{4}{x}$ ways, where x can be 0, 1, 2, 3, or 4. Thus, the probability distribution f(x) = P(X = x) is

$$f(x) = \frac{1}{16} {4 \choose x}$$
, for $x = 0, 1, 2, 3, 4$.

There are many problems where we may wish to compute the probability that the observed value of a random variable X will be less than or equal to some real number x. Writing $F(x) = P(X \le x)$ for every real number x, we define F(x) to be the cumulative distribution function of the random variable X.

$$P(x=0) = \frac{N}{N} = \frac{\binom{4}{0}}{16} = \frac{1}{16}$$

$$P(x=1) = \frac{\binom{4}{1}}{16}$$

$$P(x=2) = \frac{\binom{4}{1}}{16}$$

$$P(x=2) = \frac{\binom{4}{1}}{16}$$

$$P(x=3) = \frac{\binom{4}{0}}{16}$$

$$P(x=2) = \frac{\binom{4}{1}}{16}$$

$$P(x=1) = \frac{\binom{4}{1}}{16}$$

$$P(x=2) = \frac{\binom{4}{1}}{16}$$

$$P(x=1) = \frac{\binom{4}{1}}{16}$$

$$P(x=1) = \frac{\binom{4}{1}}{16}$$

$$P(x=2) = \frac{\binom{4}{1}}{16}$$

$$P(x=1) = \frac{\binom{4}{1}}{1$$

Definition 3.5: The cumulative distribution function F(x) of a discrete random variable X with probability distribution f(x) is

$$F(x) = P(X \le x) = \sum_{t \le x} f(t), \quad \text{for } -\infty < x < \infty.$$

For the random variable M, the number of correct matches in Example 3.2, we have

$$F(2) = P(M \le 2) = f(0) + f(1) = \frac{1}{3} + \frac{1}{2} = \frac{5}{6}$$

The cumulative distribution function of M is

The cumulative distribution function of M is Varibour Varq able Sample Space $1) | Varibour Varq able Sample Space | SJB |
<math display="block"> \frac{1}{3}, \text{ for } 0 \leq m < 1, \quad \text{ for } 0 \leq m < 1, \quad \text{ SBJ}$ $\frac{5}{6}, \text{ for } 1 \leq m < 3, \quad \text{ SBJ}$ $\frac{5}{6}, \text{ for } 1 \leq m < 3, \quad \text{ SBJ}$ $\frac{1}{3} \text{ SBJ}$ $\frac{5}{6}, \text{ for } 1 \leq m < 3, \quad \text{ SBJ}$ $\frac{5}{6}, \text{ for } 1 \leq m < 3, \quad \text{ SBJ}$ $\frac{5}{6}, \text{ SBJ}$ **JSB** JBS

Example 3.10: Find the cumulative distribution function of the random variable X in Eq. 3.9. Using F(x), verify that f(2) = 3/8.

Solution: Direct calculations of the probability distribution of Example 3.9 give f(0) = 1/16, f(1) = 1/4, f(2) = 3/8, f(3) = 1/4, and f(4) = 1/16. Therefore,

$$F(0) = f(0) = \frac{1}{16},$$

$$F(1) = f(0) + f(1) = \frac{5}{16},$$

$$F(2) = f(0) + f(1) + f(2) = \frac{11}{16},$$

$$F(3) = f(0) + f(1) + f(2) + f(3) = \frac{15}{16},$$

$$F(4) = f(0) + f(1) + f(2) + f(3) + f(4) = 1.$$
Hence,
$$\begin{cases}
0, & \text{for } x < 0, \\
\frac{1}{16}, & \text{for } 0 \le x < 1, \\
\frac{1}{16}, & \text{for } 0 \le x < 1, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1}{16}, & \text{for } 1 \le x < 2, \\
\frac{1$$




Figure 3.1: Probability mass function plot.

Figure 3.2: Probability histogram.




Figure 3.3: Discrete cumulative distribution function.

EX. 3.13: The probability distribution of A, the number of imperfections per 10 meters of a synthetic fabric in continuous rolls of uniform width, is given by:

$$F(x) = \begin{cases} 0, & \text{for } x < 0, \\ 0.41, & \text{for } 0 \le x < 1, \\ 0.78, & \text{for } 1 \le x < 2, \\ 0.94, & \text{for } 2 \le x < 3, \\ 0.99, & \text{for } 3 \le x < 4, \\ 1, & \text{for } x \ge 4. \end{cases}$$

13

• 3.2 An overseas shipment of 5 foreign automobiles contains 2 that have slight paint blemishes. If an agency receives 3 of these automobiles at random, list the elements of the sample space S using the letters B and N for blemished and nonblemished, respectively; then to each sample point assign a value x of the random variable X representing the number of automobiles purchased by the agency with paint blemishes.

Sample Space	2
NNN	0 random
NNB	1 Vorrable
NBN	1
BNN	1
NBB	2
BNB	2
BBN	2
BBB	3

3.3 Let W be a random variable giving the number of heads minus the number of tails in three tosses of a coin. List the elements of the sample space S for the three tosses of the coin and to each sample point assign a value w of W.

$$W = HHH = 3-0$$

 $W = HHT = 2-1=1$

Sample Space	w	
ПИИ	3 -	
1111T 2-1	1	
HT11 2-1	1 -	
TIIII 2-1	1-	-> 8= N
HTT 1-2	-1-	すりた
THT 1-2	-1	3,3
TT11 t-2	-1-	
TTT 0-3	-3-	





Figure 3.4: Typical density functions.

$$P(a < X < b) = \int_a^b f(x) \ dx.$$



Talong line hope but

Figure 3.5: P(a < X < b).

Definition 3.6: The function
$$f(x)$$
 is a probability density function (pdf) for the continuous random variable X , defined over the set of real numbers, if

1. $f(x) \ge 0$, for all $x \in R$.

2. $\int_{-\infty}^{\infty} f(x) dx = 1$. $\longrightarrow \ge 0 = 1$

3. $P(a < X < b) = \int_{a}^{b} f(x) dx$.

Example 3.11: Suppose that the error in the reaction temperature, in °C, for a controlled laboratory experiment is a continuous random variable X having the probability density function R = 1

$$f(x) = \begin{cases} \frac{x^2}{3}, & -1 < x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Verify that f(x) is a density function.
- (b) Find $P(0 < X \le 1)$.

Solution: We use Definition 3.6.

(a) Obviously, $f(x) \ge 0$. To verify condition 2 in Definition 3.6, we have

$$\int_{-\infty}^{\infty} f(x) \ dx = \int_{-1}^{2} \frac{x^{2}}{3} dx = \frac{x^{3}}{9} \Big|_{-1}^{2} = \frac{8}{9} + \frac{1}{9} = 1.$$

(b) Using formula 3 in Definition 3.6, we obtain

19

$$P(0 < X \le 1) = \int_0^1 \frac{x^2}{3} dx = \left. \frac{x^3}{9} \right|_0^1 = \frac{1}{9}.$$

Definition 3.7: The cumulative distribution function F(x) of a continuous random variable X with density function f(x) is

$$F(x) = P(X \le x) = \int_{-\infty}^{x} f(t) dt$$
, for $-\infty < x < \infty$.

As an immediate consequence of Definition 3.7, one can write the two results

$$P(a < X < b) = F(b) - F(a) \text{ and } f(x) = \frac{dF(x)}{dx}.$$

if the derivative exists

example 3.12: For the density function of Example 3.11, find F(x), and use it to evaluate

$$P(0 < X \le 1).$$

Solution: For
$$-1 < x < 2$$

$$F(x) = \int_{-\infty}^{x} f(t) dt = \int_{-1}^{x} \frac{x^{2}}{3} dx = \frac{x^{3}}{9} \Big|_{-1}^{x} = \frac{x^{3} + 1}{9},$$

Therefore.

- 192 in 2, seed
$$x$$
 (i. $x < -1$, $f(x) = \begin{cases} 0, & x < -1, \\ \frac{x^2+1}{9}, & -1 \le x < 2, \\ 1, & x \ge 2. \end{cases}$

The cumulative distribution function F(x) is expressed in Figure 3.6. Now

$$P(0 < X \le 1) = F(1) - F(0) = \frac{2}{9} - \frac{1}{9} = \frac{1}{9},$$

which agrees with the result obtained by using the density function in Example 3.11.




Figure 3.6: Continuous cumulative distribution function.

Example 3.13: The Department of Energy (DOE) puts projects out on bid and generally estimates what a reasonable bid should be. Call the estimate b. The DOE has determined that the density function of the winning (low) bid is

Johns
$$f(y) = \begin{cases} \frac{5}{8b}, & \frac{2}{5}b \le y \le 2b, \\ 0, & \text{elsewhere.} \end{cases}$$

Find F(y) and use it to determine the probability that the winning bid is less than the DOE's preliminary estimate b.

Solution: For $2b/5 \le y \le 2b$. $y = \int_{2b/5}^{y} \frac{5}{8b} dy = \frac{5t}{8b} \Big|_{2b/5}^{y} = \frac{5y}{8b} - \frac{1}{4}$.

Thus.

$$F(y) = \begin{cases} 0, & y < \frac{2}{5}b, \\ \frac{5y}{8b} - \frac{1}{4}, & \frac{2}{5}b \le y < 2b, \\ 1, & y \ge 2b. \end{cases}$$

To determine the probability that the winning bid is less than the preliminary bid estimate b, we have

 $P(Y \le b) = F(b) = \frac{5}{8} - \frac{1}{4} = \frac{3}{8}.$

Y ≤ b

3.6 The shelf life, in days, for bottles of a certain prescribed medicine is a random variable having the density function

$$f(x) = \begin{cases} \frac{20,000}{(x+100)^3}, & x > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the probability that a bottle of this medicine will have a shell life of

(a) at least 200 days; $\chi \ge 200$ \longrightarrow $\int_{20000} (\chi + 100)^3 dx$ (b) anywhere from 80 to 120 days

(b) anywhere from 80 to 120 days.
$$2000 (x+100)^2$$

(a) = 1/9 (b) = 0.1020

(b)= 0.1020 (c) 20000 (X+100) dx

3.7 The total number of hours, measured in units of 100 hours, that a family runs a vacuum cleaner over a period of one year is a continuous random variable X that has the density function

$$f(x) = \begin{cases} x, & 0 < x < 1, \\ 2 - x, & 1 \le x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the probability that over a period of one year, a family runs their vacuum cleaner

family runs their vacuum cleaner

(a) less than (120 hours);
$$\alpha = 120 -$$

b)
$$\int x dx \rightarrow \frac{1}{2} x \int_{0.5}^{1} (a) 0.68$$

rankom lesy.

dis

Definition 3.8:

The function f(x,y) is a joint probability distribution or probability mass function of the discrete random variables X and Y if

1.
$$f(x,y) \ge 0$$
 for all (x,y)

1.
$$f(x,y) \ge 0$$
 for all (x,y) ,
2. $\sum_{x} \sum_{y} f(x,y) = 1$,

3.
$$P(X = x, Y = y) = f(x, y)$$
.

For any region A in the xy plane, $P[(X,Y) \in A] = \sum_{A} \sum_{A} f(x,y)$.

Example 3.14: Two ballpoint pens are selected at random from a box that contains 3 blue pens. 2 red-pens, and 3 green pens. If X is the number of blue pens selected and Y is the number of red pens selected, find


(a) the joint probability function f(x, y),

$$n = n$$

(b)
$$P[(X,Y) \in A]$$
, where A is the region $\{(x,y)|x+y \le 1\}$.

Solution: The possible pairs of values (x, y) are (0, 0), (0, 1), (1, 0), (1, 1), (0, 2), and (2, 0),

(a) Now, f(0, 1), for example, represents the probability that a red and a green pens are selected. The total number of equally likely ways of selecting any 2 pens from the 8 is $\binom{5}{2} = 28$. The number of ways of selecting 1 red from 2 red pens and 1 green from 3 green pens is $\binom{2}{i}\binom{3}{i} = 6$. Hence, f(0,1) = 6/28= 3/14. Similar calculations yield the probabilities for the other cases, which are presented in Table 3.1. Note that the probabilities sum to 1. In Chapter



F(010) P(x=0 > y=0)

5, it will become clear that the joint probability distribution of Table 3.1 can be represented by the formula

$$f(x,y) = \frac{\binom{3}{x}\binom{2}{y}\binom{3}{2-x-y}}{\binom{5}{2}},$$

for x = 0, 1, 2; y = 0, 1, 2; and $0 \le x + y \le 2$.

(b) The probability that (X, Y) fall in the region A is التي مجود التي مجود عنه التي مجود عنه التي مجود التي من التي

$$P[(X,Y) \in A] = P(X+Y \le 1) = f(0,0) + f(0,1) + f(1,0),$$
$$= \frac{3}{28} + \frac{3}{14} + \frac{9}{28} = \frac{9}{14}.$$

Table 3.1: Joint Probability Distribution for Example 3.14

		f(x,y)	0	<i>x</i>	2	Row Totals		
	y	0	3 14	9 25 3 14	3 28 0	15 28 3 7	h (2d)	P(x+9 ≤4)
9 (20)		mn Totals	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	0 15 28	() 3 28	立 1		ا = ا ال دولاعمل حدوا

When X and Y are continuous random variables, the joint density function f(x,y) is a surface lying above the xy plane, and $P[(X,Y) \in A]$, where A is any region in the ry plane, is equal to the volume of the right cylinder bounded by the base A and the surface.

25

Continous

Definition 3.9: The function f(x,y) is a joint density function of the continuous random variables X and Y if

- 1. $f(x,y) \ge 0$, for all (x,y),
- $2. \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \ dx \ dy = 1$
 - 3. $P[(X,Y) \in A] = \iint_A f(x,y) dx dy$, for any region A in the xy plane.

Example 3.15: A privately owned business operates both a drive-in facility and a walk-in facility. On a randomly selected day, let X and Y, respectively, be the proportions of the time that the drive-in and the walk-in facilities are in use, and suppose that the joint density function of these random variables is

$$f(x,y) = \begin{cases} \frac{2}{5}(2x+3y), & 0 \leq x \leq 1, 0 \leq y \leq 1, \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Verify condition 2 of Definition 3.9.
- (b) Find $P[(X,Y) \in A]$, where $A = \{(x,y) \mid 0 < x < \frac{1}{2}, \frac{1}{4} < y < \frac{1}{2}\}$.

Solution: (a) The integration of f(x, y) over the whole region is

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x,y) \, dx \, dy = \int_{0}^{1} \int_{0}^{1} \frac{2}{5} (2x + 3y) \, dx \, dy$$

$$= \int_{0}^{1} \left(\frac{2x^{2}}{5} + \frac{6xy}{5} \right) \Big|_{x=0}^{x=1} dy$$

$$= \int_{0}^{1} \left(\frac{2}{5} + \frac{6y}{5} \right) dy = \left(\frac{2y}{5} + \frac{3y^{2}}{5} \right) \Big|_{0}^{1} = \frac{2}{5} + \frac{3}{5} = 1. \quad \text{The sum of } 1 = \frac{1}{5} = \frac$$

(b) To calculate the probability, we use

$$\begin{split} P[(X,Y) \in A] &= P\left(0 < X < \frac{1}{2}, \frac{1}{4} < Y < \frac{1}{2}\right) \\ &= \int_{1/4}^{1/2} \int_{0}^{1/2} \frac{2}{5} (2x + 3y) \ dx \ dy \\ &= \overline{\int_{1/4}^{1/2} \left(\frac{2x^2}{5} + \frac{6xy}{5}\right) \Big|_{x=0}^{x=1/2} \ dy} = \int_{1/4}^{1/2} \left(\frac{1}{10} + \frac{3y}{5}\right) dy \\ &= \left(\frac{y}{10} + \frac{3y^2}{10}\right) \Big|_{1/4}^{1/2} \\ &= \frac{1}{10} \left[\left(\frac{1}{2} + \frac{3}{4}\right) - \left(\frac{1}{4} + \frac{3}{16}\right)\right] = \frac{13}{160}. \end{split}$$

27

Johns along Joint Probability distirbution

Given the joint probability distribution f(x,y) of the discrete random variables X and Y, the probability distribution g(x) of X alone is obtained by summing f(x,y) over the values of Y. Similarly, the probability distribution h(y) of Y alone is obtained by summing f(x,y) over the values of X. We define g(x) and h(y) to be the marginal distributions of X and Y, respectively. When X and Y are continuous random variables, summations are replaced by integrals. We can now make the following general definition.

Definition 3.10:

The marginal distributions of
$$X$$
 alone and of Y alone are
$$g(x) = \sum_{y} f(x, y) \quad \text{and} \quad h(y) = \sum_{x} f(x, y)$$

for the discrete case, and

$$g(x) = \int_{-\infty}^{\infty} f(x, y) \, \underline{dy} \, \operatorname{and} \left(\underline{h(y)} = \int_{-\infty}^{\infty} f(x, y) \, \underline{dx} \right)$$

for the continuous case.

The term marginal is used here because, in the discrete case, the values of g(x) and h(y) are just the marginal totals of the respective columns and rows when the values of f(x, y) are displayed in a rectangular table.

Parample 3.16: Show that the column and row totals of Table 3.1 give the marginal distribution of X alone and of Y alone.

Solution: For the random variable X, we see that

$$g(0) = f(0,0) + f(0,1) + f(0,2) = \frac{3}{28} + \frac{3}{14} + \frac{1}{28} = \frac{5}{14},$$

$$g(1) = f(1,0) + f(1,1) + f(1,2) = \frac{9}{28} + \frac{3}{14} + 0 = \frac{15}{28},$$

and

$$g(2) = f(2,0) + f(2,1) + f(2,2) = \frac{3}{28} + 0 + 0 = \frac{3}{28}$$

which are just the column totals of Table 3.1. In a similar manner we could show that the values of h(y) are given by the row totals. In tabular form, these marginal distributions may be written as follows:

Table 3.

table 11 CE- (2 Summation) 1 U,N

1. Joint	Probabili	ty Dis	SUTIDI	mon	for Exam	ipie s.ra
			I		Row	
	f(x,y)	0	1	2	Totals	
	0	3	9	3 25	15 28 3	-> h(0)
y	-1	3 3 14	78 5 14	0	3 -	-> h(1) - h(9)
ju 34.	2	$\frac{1}{28}$	0	0	$\frac{1}{28}$ -	-> h(2)
Colum	nn Totals	5 14	$\frac{15}{28}$	3 25	1	110 110
		1	-7	1		Toint 1 2 / I thios.
900	1 -	96	311	1 4	1-3	الفيرة إنفع محانوا لمماول وفعلام
				9	(3)	12000
		-				(yis x 31)

navgin Vapatio

Example 3.17: Find g(x) and h(y) for the joint density function of Example 3.15. $f(x,y) = \begin{cases} \frac{2}{5}(2x+3y), & 0 \le x \le 1, 0 \le y \le 1, \\ 0, & \text{elsewhere.} \end{cases}$

Solution: By definition, $g(x) = \int_{-\infty}^{\infty} f(x,y) \, dy = \int_{0}^{1} \frac{2}{5} (2x+3y) \, dy = \left(\frac{4xy}{5} + \frac{6y^2}{10}\right)\Big|_{y=0}^{y=1} = \frac{4x+3}{5}.$ $(1) y = \frac{3}{5} + \frac{3}{5}$

$$h(y) = \int_{-\infty}^{\infty} f(x,y) \ dx = \int_{0}^{1} \frac{2}{5} (2x + 3y) \ \underline{dx} = \frac{2(1 + 3y)}{5}.$$

The fact that the marginal distributions g(x) and h(y) are indeed the probability distributions of the individual variables X and Y alone can be verified by showing that the conditions of Definition 3.4 or Definition 3.6 are satisfied. For example, in the continuous case

$$\int_{-\infty}^{\infty} g(x) \ dx = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \ dy \ dx = 1.$$

and

$$P(a < X < b) = P(a < X < b, -\infty < Y < \infty)$$

= $\int_{a}^{b} \int_{a}^{\infty} f(x, y) dy dx = \int_{a}^{b} g(x) dx$.

In Section 3.1, we stated that the value x of the random variable X represents an event that is a subset of the sample space. If we use the definition of conditional probability as stated in Chapter 2,

$$P(B|A) = \frac{P(A \cap B)}{P(A)}, \text{ provided } P(A) > 0,$$

$$P(X \mid Y) \Rightarrow \text{ leads}$$

where A and B are now the events defined by X = x and Y = y, respectively

The first where X and Y are discrete.

The first first

It is not difficult to show that the function f(x,y)/g(x), which is strictly a function of y with x fixed, satisfies all the conditions of a probability distribution. This is also true when f(x, y) and g(x) are the joint density and marginal distribution, respectively, of continuous random variables. As a result, it is extremely important that we make use of the special type of distribution of the form f(x,y)/g(x) in order to be able to effectively compute conditional probabilities. This type of distribution is called a conditional probability distribution; the formal definition follows.

Definition 3.11:

Let X and Y be two random variables, discrete or continuous. The conditional distribution of the random variable Y given that X = x is

$$f(y|x) = \frac{f(x,y)}{g(x)}, \text{ provided } g(x) > 0.$$

Similarly, the conditional distribution of X given that Y = y is

$$f(x|y) = \frac{f(x,y)}{h(y)}$$
, provided $h(y) > 0$.

31

If we wish to find the probability that the discrete random variable X falls between a and b when it is known that the discrete variable Y = y, we evaluate

$$P(a < X < b \mid Y = y) = \sum_{a < x < b} f(x|y).$$

where the summation extends over all values of X between a and b. When X and Y are continuous, we evaluate

$$P(a < X < b \mid Y = y) = \int_{a}^{b} f(x|y) dx,$$

Example 3.18: Referring to Example 3.14, find the conditional distribution of X, given that Y = 1, and use it to determine $P(X = 0 \mid Y = 1)$

Solution: We need to find f(x|y), where y = 1. First, we find that

$$h(1) = \sum_{x=0}^{2} f(x,1) = \frac{3}{14} + \frac{3}{14} + 0 = \frac{3}{7}$$

Now

$$f(x|1) = \frac{f(x,1)}{h(1)} = \left(\frac{7}{3}\right)f(x,1), \quad x = 0,1,2.$$

$$\frac{3}{19} = \underbrace{\int_{f(0,1)}^{7} f(0,1) = \left(\frac{7}{3}\right) f(0,1) = \left(\frac{7}{3}\right) \left(\frac{3}{14}\right) = \frac{1}{2}}_{f(1|1)} f(1|1) = \left(\frac{7}{3}\right) \left(\frac{3}{14}\right) = \frac{1}{2}.$$

$$f(2|1) = \left(\frac{7}{3}\right) f(2,1) = \left(\frac{7}{3}\right) (0) = 0.$$

$$f'(X = 0 \mid Y = 1) = f(0|1) = \frac{1}{2}.$$

Therefore, if it is known that 1 of the 2 pen refit probability equal to 1/2 that the other refill is not 1

			1		Row
	f(x,y)	0	1	2.	Totals
	0	MIN	9 175	Total and	15
¥	1	See les	1	0.	i
	1	101-2	i	0	1 2
		-			,
Ce	uma Totak	10	2	1	1

Table 3.1: Joint Probability Distribution for Example 3.14

Example 3.19:1 The joint density for the random variables (X,Y), where X is the unit temperature change and Y is the proportion of spectrum shift that a certain atomic particle produces, is

 $f(x,y) = \begin{cases} 10xy^2, & 0 < x < y < 1, \\ 0, & \text{elsewhere.} \end{cases}$

- (a) Find the marginal densities g(x), h(y), and the conditional density f(y|x).
- (b) Find the probability that the spectrum shifts more than half of the total observations, given that the temperature is increased by 0.25 unit.

Solution: (a) By definition,

$$\mathcal{X} = \int_{-\infty}^{\infty} f(x,y) \, dy = \int_{x}^{1} 10xy^{2} \, dy$$

$$= \frac{10}{3}xy^{3}\Big|_{y=x}^{y=1} = \frac{10}{3}x(1-x^{3}), 0 < x < 1.$$

$$h(y) = \int_{-\infty}^{\infty} f(x,y) \, dx = \int_{0}^{y} 10xy^{2} \, dx = 5x^{2}y^{2}\Big|_{x=0}^{x=y} = 5y^{4} \underbrace{0 < y < 1.}$$

Conditional Jan 8 (1 function (b) Therefore,

$$f(y|x) = \frac{f(x,y)}{g(x)} = \frac{10xy^2}{\frac{10}{3}x(1-x^3)} = \frac{3y^2}{1-x^3}, \ 0 < x < y < 1.$$

5) /2041 Mis () أيحامل حسب المعرّد المعرد المع

$$P\left(Y > \frac{1}{2} \mid X = 0.25\right) = \int_{1/2}^{1} f(y \mid x = 0.25) \ dy = \int_{1/2}^{1} \frac{3y^{2}}{1 - 0.25^{3}} \ dy = \frac{8}{9}.$$

Function 1 (8)

58,02941

Example 3.20: Given the joint density function

$$f(x,y) = \begin{cases} \frac{x(1+3y^2)}{4}, & 0 < x < 2, \ 0 < y < 1, \\ 0, & \text{elsewhere,} \end{cases}$$

find g(x), h(y), f(x|y), and evaluate $P(\frac{1}{4} < X < \frac{1}{2} \mid Y = \frac{1}{3})$. Solution: By definition of the marginal density. for 0 < x < 2,

$$\underbrace{g(x)}_{-\infty} = \int_{-\infty}^{\infty} f(x, y) \ dy = \int_{0}^{1} \frac{x(1 + 3y^{2})}{4} dy$$
$$= \left(\frac{xy}{4} + \frac{xy^{3}}{4}\right)\Big|_{y=0}^{y=1} = \frac{x}{2},$$

and for 0 < y < 1,

به عن إلعو دعل از الال 01220 C+ 477 X y 2 wilds Julia Indefendant

$$h(y) = \int_{-\infty}^{\infty} f(x, y) dx = \int_{0}^{2} \frac{x(1 + 3y^{2})}{4} dx$$
$$= \left(\frac{x^{2}}{8} + \frac{3x^{2}y^{2}}{8}\right)\Big|_{x=0}^{x=2} = \frac{1 + 3y^{2}}{2}.$$

Therefore, using the conditional density definition, for 0 < x < 2

$$f(x|y) = \frac{f(x,y)}{h(y)} = \frac{x(1+3y^2)/4}{(1+3y^2)/2} = \frac{x}{2},$$

and

$$P\left(\frac{1}{4} < X < \frac{1}{2} \mid Y = \frac{1}{3}\right) = \int_{1/4}^{1/2} \frac{x}{2} dx = \frac{3}{64}.$$

Statistical Independence

If f(x|y) does not depend on y, as is the case for Example 3.20, then f(x|y) = g(x)and f(x, y) = g(x)h(y). The proof follows by substituting

$$f(x,y) = f(x|y)h(y)$$

into the marginal distribution of X. That is,

$$g(x) = \int_{-\infty}^{\infty} f(x, y) \ dy = \int_{-\infty}^{\infty} f(x|y)h(y) \ dy.$$

If f(x|y) does not depend on y, we may write

$$g(x) = f(x|y) \int_{-\infty}^{\infty} h(y) dy.$$

Now

$$\int_{-\infty}^{\infty} h(y) \ dy = 1,$$

since h(y) is the probability density function of Y. Therefore,

$$g(x) = f(x|y)$$
 and then $f(x,y) = g(x)h(y)$.

35

Definition 3.12: Let X and Y be two random variables, discrete or continuous, with joint probability distribution f(x, y) and marginal distributions g(x) and h(y), respectively. The random variables X and Y are said to be statistically independent if and only if

$$f(x,y) = g(x)h(y)$$
 Indefendent

for all (x, y) within their range.

Example 3.21: Show that the random variables of Example 3.14 are not statistically independent. Table 3.1; Joint Probability Distribution for Example 3.14 Proof: Let us consider the point (0,1). From Table 3.1 we find the three probabilities

f(0,1), g(0), and h(1) to be

$$f(0,1) = \frac{3}{14},$$

$$g(0) = \sum_{y=0}^{2} f(0,y) = \frac{3}{28} + \frac{3}{14} + \frac{1}{28} = \frac{5}{14},$$

$$h(1) = \sum_{x=0}^{2} f(x, 1) = \frac{3}{14} + \frac{3}{14} + 0 = \frac{3}{7}.$$

$$\frac{2}{14} = \frac{5}{14} \times \frac{3}{7}$$
 $f(0,1) \neq g(0)h(1),$

前丰县

Column Totals

Row

Clearly.

and therefore X and Y are not statistically independent.

All the preceding definitions concerning two random variables can be generalized to the case of n random variables. Let $f(x_1, x_2, \dots, x_n)$ be the joint probability function of the random variables X_1, X_2, \dots, X_n . The marginal distribution of X_1 . for example, is

$$g(x_1) = \sum_{x_2} \cdots \sum_{x_n} f(x_1, x_2, \dots, x_n)$$

for the discrete case, and

$$g(x_1) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f(x_1, x_2, \dots, x_n) dx_2 dx_3 \cdots dx_n$$

for the continuous case. We can now obtain joint marginal distributions such as $g(x_1, x_2)$, where

$$g(x_1,x_2) = \begin{cases} \sum\limits_{x_3} \cdots \sum\limits_{x_n} f(x_1,x_2,\ldots,x_n) & \text{(discrete case),} \\ \int\limits_{-\infty}^{\infty} \cdots \int\limits_{-\infty}^{\infty} f(x_1,x_2,\ldots,x_n) \ dx_3 \ dx_4 \cdots dx_n & \text{(continuous case).} \end{cases}$$

We could consider numerous conditional distributions. For example, the joint conditional distribution of X_1 , X_2 , and X_3 , given that $X_4 = x_4$, $X_5 = x_5$, ..., $X_n =$ rn, is written

$$f(x_1, x_2, x_3 \mid x_4, x_5, \dots, x_n) = \frac{f(x_1, x_2, \dots, x_n)}{g(x_4, x_5, \dots, x_n)}$$

where $g(x_4, x_5, \dots, x_n)$ is the joint marginal distribution of the random variables X_4, X_5, \ldots, X_n .

A generalization of Definition 3.12 leads to the following definition for the mutual statistical independence of the variables X_1, X_2, \dots, X_n

37

Independenta 12/3 (301 00) 21 00 15 land 2 min

Definition 3.13: Let X_1, X_2, \dots, X_n be n random variables, discrete or continuous, with joint probability distribution $f(x_1, x_2, ..., x_n)$ and marginal distribution $f_1(x_1), f_2(x_2), ..., f_n(x_n)$, respectively. The random variables $X_1, X_2, ..., X_n$ are said to be mutually statistically independent if and only if

$$f(x_1, x_2, \dots, x_n) = f_1(x_1) f_2(x_2) \cdots f_n(x_n)$$

for all (x_1, x_2, \dots, x_n) within their range.

Example 3.22: Suppose that the shelf life, in years, of a certain perishable food product packaged in cardboard containers is a random variable whose probability density function is given by

$$f(x) = \begin{cases} e^{-x}, & x > 0, \\ 0, & \text{elsewhere.} \end{cases}$$

Let X_1 , X_2 , and X_3 represent the shelf lives for three of these containers selected independently and find $P(X_1 < 2, 1 < X_2 < 3, X_3 > 2)$.

Solution: Since the containers were selected independently, we can assume that the random variables X_1, X_2 , and X_3 are statistically independent, having the joint probability density

$$f(x_1, x_2, x_3) = f(x_1)f(x_2)f(x_3) = \underbrace{e^{-x_1}e^{-x_2}e^{-x_3}}_{0 = 0} = e^{-x_1 - x_2 - x_3}.$$

for
$$x_1, x_2, x_3 = f(x_1)f(x_2)f(x_3) = e^{-cx_1 - cx_2} - e^{-cx_1}$$
for $x_1 > 0$, $x_2 > 0$, $x_3 > 0$, and $f(x_1, x_2, x_3) = 0$ elsewhere. Hence
$$P(X_1 < 2, 1 < X_2 < 3, X_3 > 2) = \int_2^{\infty} \int_1^3 \int_0^2 e^{-x_1 - x_2 - x_3} dx_1 dx_2 dx_3$$

$$= (1 - e^{-2})(e^{-1} - e^{-3})e^{-2} = 0.0372.$$

- Ex. 3.47
- The amount of kerosene, in thousands of liters, in a tank at the beginning of any day is a random amount Y from which a random amount X is sold during that day. Suppose that the tank is not resupplied during the day so that x < y, and assume that the joint density 1

$$f_{f(x;y)} = \begin{cases} 2, & 0 < x \le y < 1 \\ 0, & \text{elsewhere.} \end{cases}$$

- (a) Determine if X and Y are independent.
- (b) Find P($1/4 < X < 1/2 \mid Y = 3/4$).

39



(a)
$$g(x) = 2 \int_{x}^{1} dy = 2(1-x) \text{ for } 0 < x < 1;$$
 $h(y) = 2 \int_{0}^{y} dx = 2y, \text{ for } 0 < y < 1.$

$$h(y) = 2 \int_0^y dx = 2y$$
, for $0 < y < 1$.

Since
$$f(x,y) \neq g(x)h(y)$$
, X and Y are not independent.

(b)

$$f(x|y) = f(x,y)/h(y) = 1/y$$
, for $0 < x < y$.
Therefore, $P(1/4 < X < 1/2 \mid Y = 3/4) = \frac{4}{3} \int_{1/4}^{1/2} dx = \frac{1}{3}$.

Mathematical Expectation

avy/6/6

4.1 Mean of a Random Variable

In Chapter 1, we discussed the sample mean, which is the arithmetic mean of the data. Now consider the following. If two coins are tossed 16 times and X is the number of heads that occur per toss, then the values of X are 0, 1, and 2. Suppose that the experiment yields no heads, one head, and two heads a total of 4, 7, and 5 times, respectively. The average number of heads per toss of the two coins is then

Mean =
$$\frac{(0)(4) + (1)(7) + (2)(5)}{16} = 1.06.$$

This is an average value of the data and yet it is not a possible outcome of $\{0,1,2\}$. Hence, an average is not necessarily a possible outcome for the experiment. For instance, a salesman's average monthly income is not likely to be equal to any of his monthly paychecks.

Let us now restructure our computation for the average number of heads so as to have the following equivalent form:

mean =
$$(0)\left(\frac{4}{16}\right) + (1)\left(\frac{7}{16}\right) + (2)\left(\frac{5}{16}\right) = 1.06.$$

3 1 H 5 -> 2 H

Assuming that I fair coin was tossed twice, we find that the sample space for our experiment is

$$\boldsymbol{S} = \{\boldsymbol{HH}, \boldsymbol{HT}, \boldsymbol{TH}, \boldsymbol{TT}\}$$

Since the 4 sample points are all equally likely, it follows that

and
$$P(X = 0) = P(TT) = \frac{1}{4}, \quad P(X = 1) = P(TH) + P(HT) = \frac{1}{2}, = \frac{2}{4} = \frac{1}{2}$$

$$\frac{N}{N} = \frac{1}{4} = \frac{1}{4}$$

$$\frac{1}{4} + P(X = 2) = P(HH) = \frac{1}{4}.$$

where a typical element, say TH, indicates that the first toss resulted in a tail followed by a head on the second toss. Now, these probabilities are just the relative frequencies for the given events in the long run. Therefore,

expected-mean -
$$\mu = E(X) = (0)\left(\frac{1}{4}\right) + (1)\left(\frac{1}{2}\right) + (2)\left(\frac{1}{4}\right) = 1.$$
Value

Probability Yandull

Variable

Definition 4.1: Let X be a random variable with probability distribution
$$f(x)$$
. The mean, or expected value, of X is

if X is discrete, and

$$\mu = E(X) = \sum_{x} f(x)$$

$$\mu = E(X) = \int_{-\infty}^{\infty} x f(x) dx$$

$$R = \sum_{x} f(x) dx$$

$$\mu = E(X) = \int_{-\infty}^{\infty} z f(x) dx$$
 Probabilite

if X is continuous

Example 4.1: A lot containing 7 components is sampled by a quality inspector; the lot contains
4 good components and 3 defective components. A sample of 3 is taken by the
inspector. Find the expected value of the number of good components in this

Solution: Let X represent the number of good components in the sample. The probability distribution of X is

$$f(x) = \frac{\binom{4}{x}\binom{3}{3-x}}{\binom{5}{3}}, \quad \boxed{x = 0, 1, 2, 3}$$

Simple calculations yield f(0) = 1/35, f(1) = 12/35, f(2) = 18/35, and f(3) =

$$\mu = E(X) = (0)\left(\frac{1}{35}\right) + (1)\left(\frac{12}{35}\right) + (2)\left(\frac{18}{35}\right) + (3)\left(\frac{4}{35}\right) = \frac{12}{7} = 1.7.$$

Thus, if a sample of size 3 is selected at random over and over again from a lot of 4 good components and 3 defective components, it will contain, on average, 1.7 good components

Example 4.2: A salesperson for a medical device company has two appointments on a given day. At the first appointment, he believes that he has a 70% chance to make the deal, 7 19 1500 thinks he only has a 40% chance to make the deal on the other hand, he from which, if successful, he can make \$1500. What is his based on his successful, he can make \$1500. What is his from which, if successful, he can make \$1500. What is his expected commission based on his own probability belief? Assume that the appointment results are

2500 independent of each other.

Solution: First, we know that the salesperson, for the two appointments, can have 4 possible commission totals: \$0, \$1000, \$1500, and \$2500. We then need to calculate their associated probabilities. By independence, we obtain

pated probabilities. By independence, we obtain
$$f(\$0) = (1 - 0.7)(1 - 0.1) = 0.18$$
. $f(\$2500) = (0.7)(0.4) = 0.28$. $f(\$1000) = (0.7)(1 - 0.4) = 0.42$, and $f(\$1500) = (1 - 0.7)(0.4) = 0.12$. refore, the expected commission for the salesperson is

Therefore, the expected commission for the salesperson is

إذا لمل حسارة

$$\underbrace{E(X)}_{} = (\$0)(0.18) + (\$1000)(0.12) + (\$1500)(0.12) + (\$2500)(0.28)
= \$1300.$$

dis

Continous

Example 4.3: Let X be the random variable that denotes the life in hours of a certain electronic device. The probability density function is

$$f(x) = \begin{cases} \frac{20 \text{ ood}}{x^3}, & x > 100, \end{cases}$$
 Eind the expected life of this type of device. Solution: Using Definition 4.1, we have

$$\mu = E(X) = \int_{100}^{\infty} x \frac{20,000}{x^3} \ dx = \int_{100}^{\infty} \frac{20,000}{x^2} \ dx = 200.$$

Therefore, we can expect this type of device to last, on average, 200 hours. Now let us consider a new random variable g(X), which depends on X; that is, each value of g(X) is determined by the value of X. For instance, g(X) might be X^2 or 3X - 1, and whenever X assumes the value 2, g(X) assumes the value g(2). In particular, if X is a discrete random variable with probability distribution f(x), for x = -1, 0, 1, 2, and $g(X) = X^2$, then

$$P[g(X) = 0] = P(X = 0) = f(0).$$

$$P[g(X) = 1] = P(X = -1) + P(X = 1) = f(-1) + f(1).$$

$$P[g(X) = 4] = P(X = 2) = f(2).$$

and so the probability distribution of g(X) may be written

$$\begin{array}{c|cccc} g(x) & 0 & 1 & 1 \\ \hline P[g(X) = g(x)] & f(0) & f(-1) + f(1) & f(2) \end{array}$$

By the definition of the expected value of a random variable, we obtain

$$\begin{split} \mu_{g(X)} &= \underbrace{E[g(x)]} = 0 f(0) + 1[f(-1) + f(1)] + 4 f(2) \\ &= \underbrace{(-1)^2 f(-1)} + (0)^2 f(0) + (1)^2 f(1) + (2)^2 f(2) = \sum g(x) f(x). \end{split}$$

معرب العران والمعامم

Theorem 4.1: Let X be a nuclear variable with probability distribution f(x). The expected value of the random variable g(X) is

$$\underline{\mu_{g(X)}} = E[g(X)] = \sum_x g(x) f(x)$$

if X is discrete, and

$$\mu_{\theta}(x) = E[g(X)] = \int_{-\infty}^{\infty} g(x) f(x) \ dx$$

Example 4.4: Suppose that the number of cars X that pass through a car wash between 4.00 p.m. and 5.00 p.m. on any summy Friday has the following probability distribution:

Let g(X) = 2X - 1 represent the amount of money, in dollars, paid to the attendant by the manager. Find the attendant's expected earnings for this particular time

Solution: By Theorem 4.1, the attendant can expect to receive

ملوما فحان الاقتران بنضوية بالهاالطامام

$$E[g(X)] = E(2X - 1) = \sum_{x=1}^{9} (2x - 1)f(x)$$

$$= (7) \left(\frac{1}{12}\right) + (9) \left(\frac{1}{12}\right) + (11) \left(\frac{1}{4}\right) + (13) \left(\frac{1}{4}\right)$$

$$+ (15) \left(\frac{1}{6}\right) + (17) \left(\frac{1}{6}\right) = 812.67.$$

Example 4.5: Let X be a random variable with density function

$$f(x) = \begin{cases} \frac{x^2}{3}, & -1 < x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the expected value of g(X) = 4X + 3

Solution: By Theorem 4.1, we have

$$E(4X+3) = \int_{-1}^{2} \frac{(4x+3)x^{2}}{3} dx = \frac{1}{3} \int_{-1}^{2} (4x^{3} + 3x^{2}) dx = 8.$$

We shall now extend our concept of mathematical expectation to the case of two random variables X and Y with joint probability distribution f(x, y).

Definition 4.2: Let X and Y be random variables with joint probability distribution f(x, y). The mean, or expected value, of the random variable g(X, Y) is

$$\underline{\mu_{g(X,Y)}} = E[g(X,Y)] = \sum_{x} \sum_{y} g(x,y) f(x,y)$$

if X and Y are discrete, and

$$\mu_{g(X,Y)} = E[g(X,Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x,y) f(x,y) dx dy$$

if X and Y are continuous.

Frample 4.6: Let X and Y be the random variables with joint probability distribution indicated in Table 3.1 on page 96. Find the expected value of g(X,Y) = XY. The table is reprinted here for convenience.

		100	*		Row	
	f(x,y)	0	-1	2	Totals	
15	0	28	25	3 28	15 28	
2/	1	14	14	0	+	
	2	1 28	0	- 0	1/25	
Colu	unn Totals	5	15 28	28	1	

Solution: By Definition 4.2, we write

$$\begin{split} E(XY) &= \sum_{x=0}^{2} \sum_{y=0}^{2} xyf(x,y) \\ &= (0)(0)f(0,0) + (0)(1)f(0,1) \\ &+ (1)(0)f(1,0) + (1)(1)f(1,1) + (2)(0)f(2,0) \\ &= f(1,1) = \frac{3}{14}. \end{split}$$

П

Example 4.7: Find E(Y/X) for the density function

لا كيب بعد عل عبسنال طملكنيا

$$f(x,y) = \begin{cases} \frac{x(1+3y^2)}{4}, & 0 < x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

Solution: We have

$$E\left(\frac{Y}{X}\right) = \int_0^1 \int_0^2 \frac{y(1+3y^2)}{4} \ dxdy = \int_0^1 \frac{y+3y^3}{2} \ dy = \frac{5}{8}.$$

Note that if g(X, Y) = X in Definition 4.2, we have

margin

$$E(X) = \begin{cases} \sum_{x} \sum_{y} x f(x, y) = \sum_{x} x g(x) & \text{(discrete case).} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} x f(x, y) \ dy \ dx = \int_{-\infty}^{\infty} x g(x) \ dx & \text{(continuous case).} \end{cases}$$

where g(x) is the marginal distribution of X. Therefore, in calculating E(X) over a two-dimensional space, one may use either the joint probability distribution of X and Y or the marginal distribution of X. Similarly, we define

$$E(Y) = \begin{cases} \sum_{y} \sum_{x} y f(x, y) = \sum_{y} \underline{y h(y)} & \text{(discrete case),} \\ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) \ dx dy = \int_{-\infty}^{\infty} \underline{y h(y)} \ dy & \text{(continuous case),} \end{cases}$$

where h(y) is the marginal distribution of the random variable Y.

Exercises

4.23 Suppose that X and Y have the following joint probability function:

	I amount	
flx,v)	2	4
	0.10	0.15
v 3	0.20	0.30
. 3	0.10	0.15

- (a) Find the expected value of g(XY) = XY
- (b) Find ax and av-

Answer 4.23:
(a):E[g(X,Y)] = E(XY²) =
$$\sum_{x} \sum_{y} xy^{2} f(x,y)$$
 | $\sum_{y} \frac{f(x,y)}{3} = \sum_{z} \frac{f(x,y)}{3} =$

= 2*(1)^{2*} f(2,1)+2*(3)² * f(2,3)+2*(5)² * f(2,5)+4*(1)² * f(4,1)+4*(3)² * f(4,3)+4*(5)² * f(4,5)² = 35.2

(b)
$$g(1)$$
 $g(4)$
 $\mu X = E(X) = (2)(0.1+0.2+0.1) + (4)(0.15+0.3+0.15) = 3.20 \longrightarrow {\chi g(4)}$
 $\mu Y = E(Y) = (1)(0.1+0.15) + (3)(0.2+0.3) + (5)(0.1+0.15) = 3 \longrightarrow {\chi g(4)}$
 $0 - \chi S + \frac{1}{2} + \frac{1}{2}$

The mean, or expected value, of a random variable X is of special importance in statistics because it describes where the probability distribution is centered. By itself, however, the mean does not give an adequate description of the shape of the distribution. We also need to characterize the variability in the distribution. In Figure 4.1, we have the histograms of two discrete probability distributions that have the same mean, $\mu=2$, but differ considerably in variability, or the dispersion of their observations about the mean.




Figure 4.1: Distributions with equal means and unequal dispersions.

Definition 4.3. Let X be a random variable with probability distribution f(x) and mean μ . The $\sigma^2 = E[(X - \mu)^2] = \sum_x (x - \mu)^2 f(x), \quad \text{if } X \text{ is discrete, and}$

$$\sigma^2 = E[(X - \mu)^2] = \sum_x (x - \mu)^2 f(x), \quad \text{if } X \text{ is discrete, and } \int_{-\infty}^{\infty} (x - \mu)^2 f(x) \, dx, \quad \text{if } X \text{ is continuous.}$$

The positive square root of the variance, σ_s is called the standard deviation of

E(X-M) f(x)

I sample 4.8. Let the random variable X represent the number of automobiles that are used for official business purposes on any given workday. The probability distribution for company 4 [Figure L1(a)] is

Show that the variance of the probability distribution for company B is greater than that for company A. Substinu: For conquity A, we find that

$$\mu_A = L(X) = (1)(0.3) + (2)(0.4) + (3)(0.3) = 2.0,$$

East Probability $\sigma_A^2 = \sum_{i=1}^{n} (x-2)^2 = (1-2)^2(0.3) + (2-2)^2(0.4) + (3-2)^2(0.3) = 0.6.$

For company B, we have

$$\mu_B = E(X) = (0)(0.2) + (1)(0.1) + (2)(0.3) + (3)(0.3) + (4)(0.1) = 2.0,$$
 and then

$$\sigma_D^2 = \sum_{s=0}^{4} (s-2)^2 f(s)$$

$$= 2(t-2)^2 (0.3) + (1-2)^2 (0.1) + (2-2)^2 (0.1)$$

$$+ (3-2)^2 (0.3) + (3-2)^2 (0.1) = 1.6.$$

Theorem 4.2: The variance of a random variable X is

$$\sigma^2 = E(X^2) - \mu^2.$$

Proof: For the discrete case, we can write

$$\sigma^{2} = \sum_{x} (x - \mu)^{2} f(x) = \sum_{x} (x^{2} - 2\mu x + \mu^{2}) f(x)$$
$$= \sum_{x} x^{2} f(x) - 2\mu \sum_{x} x f(x) + \mu^{2} \sum_{x} f(x).$$

Since $\mu = \sum_{x} x f(x)$ by definition, and $\sum_{x} f(x) = 1$ for any discrete probability distribution, it follows that

$$\sigma^2 = \sum x^2 f(x) - \mu^2 = E(X^2) - \mu^2.$$

For the continuous case the proof is step by step the same, with summations replaced by integrations.

Example 4.9: Let the random variable X represent the number of defective parts for a machine when 3 parts are sampled from a production line and tested. The following is the probability distribution of X. Using Theorem 4.2, calculate σ^2 .

Solution: First, we compute

$$\mu = (0)(0.51) + (1)(0.38) + (2)(0.10) + (3)(0.01) = 0.61.$$

Now

$$E(X^2) = (0)(0.51) + (1)(0.38) + (4)(0.10) + (9)(0.01) = 0.87.$$

Therefore,

$$\sigma^2 = 0.87 - (0.61)^2 = 0.4979.$$

Example 4.10: The weekly demand for a dranking-water product, in thousands of liters, from a local chain of efficiency stores is a continuous random variable X having the probability density

$$f(x) = \begin{cases} 2(x-1), & 1 < x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the mean and variance of X

$$\mu = E(X) = 2\int_1^2 x(x-1) \ dx = \frac{5}{3}$$

$$E(X^2) = 2 \int_1^2 x^2 (x-1) \ dx = \frac{17}{6}$$

Therefore,

$$\sigma^2 = \frac{17}{6} - \left(\frac{5}{3}\right)^2 = \frac{1}{18}.$$

Definition 4.3: Let X be a random variable with probability distribution f(z) and mean μ . The variance of X is

$$\begin{split} \sigma^2 &= E[(X-\mu)^2] = \sum_x (x-\mu)^2 f(x), & \text{if } X \text{ is discrete, and} \\ \sigma^2 &= E[(X-\mu)^2] = \int_{-\infty}^\infty (x-\mu)^2 f(x) \ dx, & \text{if } X \text{ is containtous.} \end{split}$$

The positive square root of the variance, a, is called the standard deviation of

The variance of a random variable X is

$$\sigma^2 = E(X^2) - \mu^2.$$

Theorem 4.3: Let X be a random variable with probability distribution f(x). The variance of the random variable g(X) is

$$\sigma_{g(X)}^2=E\{[g(X)-\mu_{g(X)}]^2\}=\sum_x[g(x)-\mu_{g(X)}]^2f(x)$$
 if X is discrete, and

 $\sigma_{g(X)}^2 = E\{[g(X) - \mu_{g(X)}]^2\} = \int_{-\infty}^{\infty} [g(x) - \mu_{g(X)}]^2 f(x) \ dx$

if X is continuous.

Proof: Since g(X) is itself a random variable with mean $\mu_{g(X)}$ as defined in Theorem 4.1, it follows from Definition 4.3 that

$$\sigma_{g(X)}^2 = E\{ [g(X) - \mu_{g(X)}] \}.$$

Now, applying Theorem 4.1 again to the random variable $[g(X) - \mu_{g(X)}]^2$ completes the proof.

Example 4.11: Calculate the variance of
$$g(X) = 2X + 3$$
, where X is a random variable where $g(X) = 2X + 3$ w

$$\sigma_{g(X)}^2 = \sum_x [g(x) - \mu_{g(X)}]^2 f(x)$$

$$\mu_{2X+3} = E(2X+3) = \sum_{x=0}^{3} (2x+3)f(x) = 6.$$

Now, using Theorem 4.3, we have

$$\sigma_{2X+3}^2 = E\{[(2X+3) - \mu_{2x+3}]^2\} = E[(2X+3-6)^2]$$

$$= E(4X^2 - 12X + 9) = \sum_{x=0}^{3} (4x^2 - 12x + 9)f(x) = 4.$$

Example 4.12a Let X be a random variable having the density function given in Example 4.5 on page 115. Find the variance of the random variable Q(X) = 4X + 3.

Solution: In Example 4.5, we found that $\mu_{4X+3} = 8$. Now, using Theorem

$$\sigma_{4X+3}^2 = E\{[(4X+3)-8]^2\} = E[(4X-5)^2]$$

$$= \int_{-1}^{2} (4x-5)^2 \frac{r^2}{3} dr = \frac{1}{3} \int_{-1}^{2} (16r^4 - 40r^3 + 25r^2) dr = \frac{51}{5}.$$

If $g(X,Y) = (X - \mu_X)(Y - \mu_Y)$, where $\mu_X = E(X)$ and $\mu_Y = E(Y)$, Definition 4.2 yields an expected value called the covariance of X and Y, which we denote by σ_{XY} or Cov(X,Y).

Let X be a random variable with density function

$$f(x) = \begin{cases} \frac{x^2}{3}, & -1 < x < 2, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the expected value of g(X) = 4X + 3,

Definition 4.4: Let X and Y be random variables with joint probability distribution f(x, y). The covariance of X and Y is

$$\sigma_{XY} = E[(X - \mu_X)(Y - \mu_Y)] = \sum_x \sum_y (x - \mu_X)(y - \mu_y) f(x,y)$$

if X and Y are discrete, and

$$\sigma_{XY} = E[(X - \mu_X)(Y - \mu_Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_X)(y - \mu_y)f(x, y) \ dx \ dy$$

if X and Y are continuous

Theorem 4.4: The covariance of two random variables X and Y with means μ_X and μ_Y , respectively, is given by

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y$$

Proof: For the discrete case, we can write

$$\sigma_{XY} = \sum_{x} \sum_{y} (x - \mu_X)(y - \mu_Y) f(x, y)$$

$$= \sum_{x} \sum_{y} xy f(x, y) - \mu_X \sum_{x} \sum_{y} y f(x, y)$$

$$- \mu_Y \sum_{x} \sum_{y} x f(x, y) + \mu_X \mu_Y \sum_{x} \sum_{y} f(x, y).$$

Since

$$\mu_{\scriptscriptstyle X} = \sum_x x f(x,y), \quad \mu_{\scriptscriptstyle Y} = \sum_y y f(x,y), \text{ and } \sum_x \sum_y f(x,y) = 1$$

for any joint discrete distribution, it follows that

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y - \mu_Y \mu_X + \mu_X \mu_Y = E(XY) - \mu_X \mu_Y.$$

For the continuous case, the proof is identical with summations replaced by integrals.

 $E(XY) = \sum_{i=1}^{2} \sum_{j=1}^{2} ryf(x,y)$

 $= f(1,1) = \frac{3}{14}$

= (0)(0)f(0,0) + (0)(1)f(0,1)

+(1)(0)f(1,0) + (1)(1)f(1,1) + (2)(0)f(2,0)

Example 4.13: Example 3.14 on page 95 describes a situation involving the number of blue refills X and the number of red refills Y. Two refills for a ballpoint pen are selected at random from a certain box, and the following is the joint probability distribution:

	GREET !	1	x		
	f(x, y)	0	1	2	h(y)
3.	0	3 28	28	3 28	$\frac{15}{28}$
y	1	3	$\frac{3}{14}$	0	7
- 1	2	$\frac{1}{28}$	0	0	28
	g(x)	5	$\frac{15}{28}$	$\frac{3}{28}$	1

Find the covariance of X and Y

Solution: From Example 4.6, we see that E(XY) = 3/14. Now

$$\mu_X = \sum_{x=0}^{2} xg(x) = (0) \left(\frac{5}{14}\right) + (1) \left(\frac{15}{28}\right) + (2) \left(\frac{3}{28}\right) = \frac{3}{4}.$$

and

$$\mu_Y = \sum_{y=0}^2 y h(y) = (0) \left(\frac{15}{28}\right) + (1) \left(\frac{3}{7}\right) + (2) \left(\frac{1}{28}\right) = \frac{1}{2}.$$

Therefore

$$\sigma_{xy} = E(XY) - \mu_{x}\mu_{y} = \frac{3}{14} - \left(\frac{3}{4}\right)\left(\frac{1}{2}\right) = -\frac{9}{56}$$
(العلاقة لعد كالمالي ير ل على إن 24 حدث بنعلي كالا (العلاقة لين يرو)

$$f(x,y) = \begin{cases} 8xy, & 0 \le y \le x \le 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the covariance of X and Y.

Solution: We first compute the marginal density functions. They are

$$g(r) = \begin{cases} 4x^3, & 0 \le x \le 1, \\ 0, & \text{elsewhere.} \end{cases}$$

and

$$h(y) = \begin{cases} 4y(1-y^2), & 0 \le y \le 1, \\ 0, & \text{elsewhere.} \end{cases}$$

From these marginal density functions, we compute

$$\mu_X = E(X) = \int_0^1 4x^4 dx = \frac{4}{5} \text{ and } \mu_Y = \int_0^1 4y^2 (1 - y^2) dy = \frac{8}{15}.$$

From the joint density function given above, we have


$$E(XY) = \int_0^1 \int_y^1 8x^2y^2 \ dx \ dy = \frac{4}{9}.$$

Then

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y = \frac{4}{9} - \left(\frac{4}{5}\right) \left(\frac{8}{15}\right) = \frac{4}{225}.$$

Л

Correlation Coefficient



Definition 4.5: Let X and Y be random variables with covariance σ_{XY} and standard deviations σ_{X} and σ_{Y} , respectively. The correlation coefficient of X and Y is

$$\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$$

Praniple 4.15: Find the correlation coefficient between X and Y in Example 4.13

Solution: Since

$$E(X^2) = (0^2) \left(\frac{5}{14}\right) + (1^2) \left(\frac{15}{28}\right) + (2^2) \left(\frac{3}{28}\right) = \frac{27}{28}$$

$$E(Y^2) = (0^2) \left(\frac{15}{28}\right) + (1^2) \left(\frac{3}{7}\right) + (2^2) \left(\frac{1}{28}\right) = \frac{4}{7}.$$

$$\mu_X = \sum_{x=0}^{\infty} x g(x) = (0) \left(\frac{5}{14}\right) + (1) \left(\frac{15}{28}\right) + (2) \left(\frac{3}{28}\right) = \frac{3}{4}.$$

$$\sigma_X^2 = \frac{27}{28} - \left(\frac{3}{4}\right)^2 = \frac{45}{112}$$

$$\mu_Y = \sum_{y=0}^2 y h(y) = (0) \left(\frac{15}{28}\right) + (1) \left(\frac{3}{7}\right) + (2) \left(\frac{1}{28}\right) = \frac{1}{2}$$

$$\sigma_Y^2 = \frac{1}{7} - \left(\frac{1}{2}\right)^2 = \frac{9}{28}.$$

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y = \frac{3}{11} - \left(\frac{3}{4}\right) \left(\frac{1}{2}\right) = -\frac{9}{56}.$$

$$\mu_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y} = \frac{-9/56}{\sqrt{(15/112)(9/28)}} = -\frac{1}{\sqrt{5}}.$$
Plogability

The variance of a random variable X is

$$\sigma^2 = E(X^2) - \mu^2,$$

			T			
	f(x,y)	0	1	2	4(9)	
	0	28	75	黄	15	
W.	1	3	3	0	3	
	- 1	11	0	- 65		
	-	28	-		25	
	g(x)	-	12	20	TO S	

$$\begin{split} E(XY) &= \sum_{x=0}^{2} \sum_{y=0}^{2} xy f(x,y) \\ &= (0)(0) f(0,0) + (0)(1) f(0,1) \\ &+ (1)(0) f(1,0) + (1)(1) f(1,1) + (2)(0) f(2,0) \\ &= f(1,1) = \frac{3}{14}. \end{split}$$

Solution: Because

$$E(X^2) = \int_0^1 \underbrace{4x^5 dx} = \frac{2}{3} \text{ and } E(Y^2) = \int_0^1 4y^3 (1 - y^2) dy = 1 - \frac{2}{3} = \frac{1}{3}.$$

we conclude that

$$\sigma_X^2 = \frac{2}{3} - \left(\frac{4}{5}\right)^2 = \frac{2}{75} \text{ and } \sigma_Y^2 = \frac{1}{3} - \left(\frac{8}{15}\right)^2 = \frac{11}{225}$$

$$\sigma_{XY} = E(XY) - \mu_X \mu_Y = \frac{4}{9} - \left(\frac{4}{5}\right) \left(\frac{8}{15}\right) = \frac{1}{225}$$

$$\rho_{XY} = \frac{4/225}{\sqrt{(2/75)(11/225)}} = \frac{1}{\sqrt{66}}$$

$$\sigma^2 = E(X^2) - \mu^2$$

$$\sigma^2 = E(X^2) - \mu^2$$

$$\sigma(1 - \left[\frac{\mu^2}{\mu}, \frac{\mu_{2,2}}{\mu_{2,2}}\right]$$

$$\sigma(2 - E(X) - \left[\frac{\mu^2}{\mu}, \frac{\mu_{2,2}}{\mu_{2,2}}\right]$$

$$\sigma(3 - E(X) - \left[\frac{\mu^2}{\mu}, \frac{\mu_{2,2}}{\mu_{2,2}}\right]$$

$$\sigma(4 - E(X) - \mu^2)$$

$$\sigma(4 - E(X$$

Theorem 4.7: The expected value of the sum or difference of two or more functions of the random variables X and Y is the sum or difference of the expected values of the functions. That is,

 $Z[g], \neq S) \Leftrightarrow \mathcal{E}[g(X,Y) \pm h(X,Y)] = E[g(X,Y)] \pm E[h(X,Y)].$

Proof: By Definition 4.2,

$$\begin{split} E[g(X,Y) &\pm h(X,Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [g(x,y) \pm h(x,y)] f(x,y) \ dx \ dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x,y) f(x,y) \ dx \ dy \pm \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x,y) f(x,y) \ dx \ dy \\ &= E[g(X,Y)] \pm E[h(X,Y)]. \end{split}$$

Corollary 4.3: Setting g(X,Y) = g(X) and h(X,Y) = h(Y), we see that $E[g(X) \pm h(Y)] = E[g(X)] \pm E[h(Y)].$

 $M_{*}(X_{*}Y_{*}) = Y_{*}$ we see that

$$E[X\pm Y] = E[X] \pm E[Y]$$

If X represents the daily production of some item from machine A and Y the daily production of the same kind of item from machine B, then X+Y represents the total number of items produced daily by both machines. Corollary 4.4 states that the that the average daily production for both machines is equal to the sum of the average daily production of each machine.

Theorem 4.8: Let X and Y be two independent random variables. Then

$$E(XY) = E(X)E(Y).$$

Proof: By Definition 4.2.

$$E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x,y) dx dy.$$

п

Since X and Y are independent, we may write

$$f(x,y)=g(x)h(y), \\$$

where g(x) and h(y) are the marginal distributions of X and Y, respectively. Hence,

$$\begin{split} E(XY) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyg(x)h(y) \ dx \ dy = \int_{-\infty}^{\infty} xg(x) \ dx \int_{-\infty}^{\infty} yh(y) \ dy \\ &= E(X)E(Y), \end{split}$$

Corollary 4.5; Let X and Y be two independent random variables. Then $\sigma_{XY} = 0$,

Proof: The proof can be carried out by using Theorems 4.4 and 4.8.

$$\mathrm{Cov}(X,Y) := \mathbb{E}\left[(X - \mu_X) (Y - \mu_Y) \right]$$

$$\begin{aligned} \operatorname{Cov}(X,Y) &= \mathbb{E}\left[(X - \mu_X)(Y - \mu_Y) \right] \\ &= \mathbb{E}[(X - \mu_X)] \mathbb{E}[(Y - \mu_Y)] \\ &= (\mathbb{E}[X] - \mu_X)(\mathbb{E}[Y] - \mu_Y) \\ &= 0. \end{aligned}$$

4.31 It is known that the ratio of gallium to arsenide does not affect the functioning of gallium arsenide waters, which are the main components of microchips. Let X denote the ratio of gallium to arsenide and Y denote the functional waters retrieved during a 1-hour period. X and Y are independent random variables with the joint density function.

$$f(x,y) = \begin{cases} \frac{x(1+3y^2)}{1}, & 0 < x < 2, \ 0 < y < 1, \\ 0, & \text{elsewhere.} \end{cases}$$

Show that E(XY) = E(X)E(Y), as Theorem 4.8 suggests. Solution: By definition,

$$E(XY) = \int_0^1 \int_0^2 \frac{x^2y(1+3y^2)}{4} \ dxdy = \frac{5}{6}, \ E(X) = \frac{4}{3}, \ \text{and} \ E(Y) = \frac{5}{8}.$$

Hence,

$$E(X)E(Y) = \left(\frac{4}{3}\right)\left(\frac{5}{8}\right) = \frac{5}{6} = E(XY).$$

We conclude this section by proving one theorem and presenting several corollaries that are useful for calculating variances or standard deviations. e (Permutation) o ustalian) different de core dismiliales de Position billiques (silis) Colision avvangement &

Permutation

Circular

(n-1)!

بحون المني دانوي

يمنى وينا ماوجيت محلقة

e !! duli des circular

1) nobject iletie - wit

الخالفة في محطده بية خواع اله

2) nobject + lein Y to

npr (n-r) 3 222 21/2 .3) nobject nicht

110,3 60, cds 15 5 9 013 Jis 7! 4! - clear de cue!

4) nobject

NI Walied Millingion

n! & (combinations) (i)

Usolo postion) 2 lois, licit con + 50 Just selection

Reyword: (select / elect/ + o get)

CITCHAR

Ex.2.44

In how many ways can a caravan of 8 covered wagons from Arizona be arranged in a circle?

(n-1)! = 7!

5040

45

3 9181

Ex 2.46

In how many ways can 3 oaks, 4 pines, and 2 maples be arranged along a property line if one does not distinguish among trees of the same kind?

A college plays 12 football games during a season. In how many ways can the team end the season with 7 wins, 3 losses, and 2 ties?

Ex 2.49

How many ways are there to select 3 candidates from 8 equally qualified recent graduates for openings in an accounting firm

$$\frac{n!}{7!(n-r)!}$$
56 ways
$$\frac{8!}{8!}$$
Uscales (8) - $\frac{8!}{3!(8-3)!}$
Permatation

• 2.36 (a) How many three-digit numbers can be formed from the digits 0, 1, 2, 3, 4, 5, and 6, if respectively each digit can be used only once? (b) How many of these are odd numbers? C) Howmany are greater

180 (6+6+5)

180 (6+6+5)

180 (6+6+5)

180 (6+6+5)

180 (6+6+5)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (5+5+3)

180 (

Definition 2.9: The probability of an event A is the sum of the weights of all sample points in A. Therefore,

Solution of the second of the

points in N=40620

This DI Ja U J3 50

H

Furthermore, if A_1, A_2, A_3, \ldots is a sequence of mutually exclusive events, then

 $P(A_1 \cup A_2 \cup A_3 \cup \cdots) = P(A_1) + P(A_2) + P(A_3) + \cdots$

Example 2.24: A coin is tossed twice. What is the probability that at least 1 head occurs?

Solution: The sample space for this experiment is

 $S = \{ \underline{HH}, \underline{HT}, \underline{TH}, \underline{TT} \}.$

If the coin is balanced, each of these outcomes is equally likely to occur. Therefore, we assign a probability of ω to each sample point. Then $4\omega=1$, or $\omega=1/4$. If A represents the event of at least 1 head occurring, then

$$A = \{HH, HT, TH\}$$
 and $P(A) = \frac{1}{4} + \frac{1}{4} + \frac{1}{4} = \frac{3}{4}$.

If more than one head = 1

Example 2.25: A die is loaded in such a way that an even number is twice as likely to occur as an odd number. If E is the event that a number less than 4 occurs on a single toss of the die, find P(E).

the die, find P(E).

even = 20dd $\int_{0}^{1} \int_{0}^{2} \int_{0}^{2}$

Solution: The sample space is $S = \{1, 2, 3, 4, 5, 6\}$. We assign a probability of w to each odd number and a probability of 2w to each even number. Since the sum of the probabilities must be 1, we have 9w = 1 or w = 1/9. Hence, probabilities of 1/9 and 2/9 are assigned to each odd and even number, respectively. Therefore,

1 1213
$$< 4$$

$$E = \{1,2,3\} \text{ and } P(E) = \frac{1}{9} + \frac{2}{9} + \frac{1}{9} = \frac{4}{9}.$$

$$E = \{1,2,3\} \text{ and } P(E) = \frac{1}{9} + \frac{2}{9} + \frac{1}{9} = \frac{4}{9}.$$

 $E \rightarrow more than 5$ E = 263 = 29 $2w^2 = 9$

511213/4/5/63 2W 2W 2W 2W A= 52, 4,63 PA= 5

Example 2.26: In Example 2.25, let A be the event that an even number turns up and let B be solution: For the events $A = \{2, 4, 6\}$ and $B = \{3, 6\}$, we have

By assigning a probability of 1/9 to each odd number and 2/9 to each even number, $P_{B} = 9 = \frac{1}{9} = \frac{1}{9}$ we have

$$P(A \cup B) = \frac{2}{9} + \frac{1}{9} + \frac{2}{9} + \frac{2}{9} = \frac{7}{9} \text{ and } P(A \cap B) = \frac{2}{9}.$$

If the sample space for an experiment contains N elements, all of which are equally likely to occur, we assign a probability equal to 1/N to each of the N points. The probability of any event A containing n of these N sample points is then the ratio of the number of elements in A to the number of elements in S.

ι

53

$$P(A) = \frac{n}{N}$$
.

Example 2.27: A statistics class for engineers consists of 25 industrial, 10 mechanical, 10 electrical, and 8 civil engineering students. If a person is randomly selected by the instructor to answer a question, find the probability that the student chosen is (a) an industrial engineering major and (b) a civil engineering or an electrical engineering major.

M = 10

Solution: Denote by I, M, E, and C the students majoring in industrial, mechanical, electrical, and civil engineering, respectively. The total number of students in the class is 53, all of whom are equally likely to be selected.

(a) Since 25 of the 53 students are majoring in industrial engineering, the probability of event I, selecting an industrial engineering major at random, is

 $P(I) = \frac{25}{53}.$

(b) Since 18 of the 53 students are civil or electrical engineering majors, it follows ICU IEK

$$P(C \cup E) = \frac{18}{53} = \frac{8+10}{53} = \frac{18}{53}$$

P(C)+P(E)-P(CNE)= = 8+10-0

if do joint every Theorem 2.7: If A and B are two events, then $P(A \cup B) = P(A) + P(B) - P(A \cap B).$




Figure 2.7: Additive rule of probability.

11,000

Example:

John is going to graduate from an industrial engineering department in a university by the end of the semester. Alter being interviewed at two companies he likes, he assesses that his probability of getting an offer from PA) = 0.8 P(B) = 06 company A is 0.8, and the probability that he gets an offer from company B is 0.6. If on the other hand, he believes that the probability that he will get offers from P(AAB)=0 Both companies is 0.5, what is the probability that he will get at least one offer from these two companies?

 $P(A \cup B) = P(A) + P(B) - P(A f \mid B) = 0.8 + 0.6 - 0.5 = 0.9.$

55

Corollary 2.1: If A and B are mutually exclusive, then

$$P(A \cup B) = P(A) + P(B)$$
. disjoint

Corollary 2.1 is an immediate result of Theorem 2.7, since if A and B are mutually exclusive, $A \cap B = 0$ and then $P(A \cap B) = P(\phi) = 0$. In general, we can write Corollary 2.2.

Corollary 2.2: If
$$A_1, A_2, \ldots, A_n$$
 are mutually exclusive, then

$$P(A_1 \cup A_2 \cup \cdots \cup A_n) = P(A_1) + P(A_2) + \cdots + P(A_n).$$


A collection of events $\{A_1, A_2, \dots, A_n\}$ of a sample space S is called a partition of S if A_1, A_2, \ldots, A_n are mutually exclusive and $A_1 \cup A_2 \cup \cdots \cup A_n = S$. Thus, we have

Corollary 2.3: If
$$A_1, A_2, \dots, A_n$$
 is a partition of sample space S , then
$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n) = P(S) = 1.$$

As one might expect, Theorem 2.7 extends in an analogous fashion.

Theorem 2.8: For three events
$$A$$
, B , and C ,

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$
$$-P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C).$$



Solution: Let G, W, R, and B be the events that a buyer selects, respectively, a green, white, red, or blue automobile. Since these four events are mutually exclusive, the probability is

 $P(\underline{G} \cup \underline{W} \cup \underline{R} \cup \underline{B}) = P(G) + P(W) + P(R) + P(B)$ = 0.09 + 0.15 + 0.21 + 0.23 = 0.68

11-AB

Often it is more difficult to calculate the probability that an event occurs than it is to calculate the probability that the event does not occur. Should this be the case for some event A, we simply find P(A') first and then, using Theorem 2.7, find P(A) by subtraction.

Theorem 2.9: If A and A' are complementary events, then P(A) + P(A') = 1.A={21416} **Proof:** Since $A \cup A' = S$ and the sets A and A' are disjoint, $1 = P(S) = P(A \cup A') = P(A) + P(A').$ $A' = (1 - \frac{3}{2})$

Example 2.32: If the probabilities that an automobile mechanic will service 3, 4, 5, 6, 7, or 8 or more cars on any given workday are, respectively, 0.12, 0.19, 0.28, 0.24, 0.10, and 8 0.07, what is the probability that he will service at least 5 cars on his next day at work?

P(E) = P(5) + P(6) + P(7) + P(8)P(E)= 0.28+0.24+0.1+0.07 P(E) = 0.69

Let E be the event that at least 5 cars are serviced. Now, P(E) = 1 - P(E')where E' is the event that fewer than 5 cars are serviced. Since 1- P(3)-P(4)

$$P(E') = 0.12 + 0.19 = 0.31,$$

it follows from Theorem 2.9 that

P(E) = 1 - 0.31 = 0.69.

2.58 An automobile manufacturer is concerned about a possible recall of its best-selling four-door sedan. If there were a recall, there is 0.25 probability that a defect is in the brake system, 0.18 in the transmission, 0.17 in the fuel system, and 0.40 in some other area.

(a) What is the probability that the defect is the brakes U or the fueling system if the probability of defects in both systems simultaneously is 0.15?

(b) What is the probability that there are no defects conferent in either the brakes or the fueling system?

the probability that there he probability that there he brakes or the fueling system?

(a) = 0.27
$$\rightarrow P(B) \rightarrow P(F) - P(B \cap F)$$

(b) = 0.73

$$P(B' \cap F') = 1 - P(B \cup F)$$

$$= 1 - 0.27$$

[1990-2010]

PI

Example 2.33: Suppose the manufacturer's specifications for the length of a certain type of computer cable are 2000 ± 10 millimeters. In this industry, it is known that small cable 0,005 is just as likely to be defective (not meeting specifications) as large cable. That is

1-0.99 =000

1

the probability of randomly producing a cable with length exceeding 2010 millimeters is equal to the probability of producing a cable with length smaller than 1990 millimeters. The probability that the production procedure meets specifications is Objects 4 d=q known to be 0.99.

(a) What is the probability that a cable selected randomly is too large?

(b) What is the probability that a randomly selected cable is larger than 1990 millimeters?

Solution: Let M be the event that a cable meets specifications. Let S and L be the events that the cable is too small and too large, respectively. Then

(a)
$$P(M) = 0.99$$
 and $P(S) = P(L) = (1 - 0.99)/2 = 0.005$.

(b) Denoting by X the length of a randomly selected cable, we have


$$P(1990 \le X \le 2010) = P(M) = 0.99.$$

Since
$$P(X \ge 2010) = P(L) = 0.005$$
, $0.99 + 0.005$
 $1990 \text{ Y} P(X \ge 1990) = P(M) + P(L) = 0.995$.

This also can be solved by using Theorem 2.9:

$$P(X \ge 1990) + P(X < 1990) = 1.$$

Thus,
$$P(X \ge 1990) = 1 - P(S) = 1 - 0.005 = 0.995$$
.



Conditional Probability

Question:

In a group of 100 sports car buyers, 40 bought alarm systems, 30 purchased bucket seats, and 20 purchased an alarm system and bucket seats. If a car buyer chosen at random bought an alarm system, what is the probability they also bought

alarm bucket seats?

P(B/A)

P(A) = $\frac{40}{100}$ Bucket Seats

P(A) B

P(A) = $\frac{40}{100}$ P(A) Bucket Seats

Answer:

 $P(B|A) = P(A \cap B) / P(A) = 0.2 / 0.4 = 0.5$

The probability that a buyer bought bucket seats, given that they purchased an alarm system, is 50%

63

Example 2.34: The probability that a regularly scheduled flight departs on time is P(D) = 0.83: the probability that it arrives on time is P(A) = 0.82; and the probability that it departs and arrives on time is $P(D \cap A) = 0.78$. Find the probability that a plane

(a) arrives on time, given that it departed on time, and (b) departed on time, given that it has arrived on time.

Solution: Using Definition 2.10, we have the following.

(a) The probability that a plane arrives on time, given that it departed on time, is

$$P(A|D) = \frac{P(D \cap A)}{P(D)} = \frac{0.78}{0.83} = \underline{0.94}.$$

(b) The probability that a plane departed on time, given that it has arrived on time, is

$$P(D|A) = \frac{P(D \cap A)}{P(A)} = \frac{0.78}{0.82} = 0.95.$$

الحوال يعتود على اطقام

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being produced. These strips can be defective in two ways, length and nature of texture. For the case of the latter, the process of identification is very complicated. It is known from historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests. If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

alas 1

(31)

فيل

Solution: Consider the events

L: length defective. T: texture defective. Given that the strip is length defective, the probability that this strip is texture defective is given by

P(2) = 0,01 $P(T|L) = \frac{P(T \cap L)}{P(L)} = \frac{0.008}{0.1} = 0.08.$ P(F) = 0.5 Texture) de l'enothes s'ist given est pl d'émble & P(TAL) = 0.008

(T/1) TO:

65

Two events A and B are independent if and only if P(A) & P(B)

Definition 2.11:

P(B|A) = P(B) or P(A|B) = P(A), P(A) P(B)

assuming the existences of the conditional probabilities. Otherwise, A and B are dependent.

Theorem 2.10: If in an experiment the events A and B can both occur, then

 $2 \cap P(A \cap B) = P(A)P(B|A)$ provided P(A) > 0. desendent

Example 2.36: Suppose that we have a fuse box containing 20 fuses, of which 5 are defective. If عن ورما المراجع لا عن المراجع 2 fuses are selected at random and removed from the box in succession without replacing the first, what is the probability that both fuses are defective?

Solution: We shall let A be the event that the first fuse is defective and B the event that the second fuse is defective; then we interpret $A \cap B$ as the event that A occurs and then B occurs after A has occurred. The probability of first removing a defective fuse is 1/4: then the probability of removing a second defective fuse from the remaining 4 is 4/19. Hence,

0->5 NOND-15

dependent $P(A \cap B) = \left(\frac{1}{4}\right)\left(\frac{4}{19}\right) = \frac{1}{19}. \quad P(A) P(B|A)$ P(A)= 是 = 4 P(B)A) = 4

Stide 66

$$P(R) = \frac{3}{6}$$

احتمالية ع إذا بحمالحرة حراء وعارجماع

defendent = (3121) CIXITE DE USIN CIXI 10 - Independent dependent

The grabability that and merican industry will locate in shanghair China is 0.7, the probability that it will locate in Beising, china is out and the probability that it will locate in either shanghai or Beijing or both is 0.8, what is the probability that the industry will locate (a) in both cities? b) in neither city?

a) p(sUB) = P(s)+P(B)-P(SAB) 0.8 = 0.7 to.4 - P(SAB)


P(s) = 0.7 P(B)=0.4


VSUB)=00


P(51B)=03

b) P(SNB) = 1-(SUB) = 1 - 0.8 = 0.2

NOTEBOOK







Probability that the fire engine and one ambulance available for emergencies. The that the ambulance is available when needed is 0.98, and the probability resulting from a burning building, find the probability that both the ambulance and the fire engine will be available, assuming they operate independently.

Solution: Let A and B represent the respective events that the fire engine and the ambulance are available. Then

P(A \cap B) = P(A)P(B) = (0.98)(0.92) = 0.9016.

P(B)=0.92

69

Ex. 2.95

One overnight case contains 2 bottles of aspirin and 3 bottles of thyroid tablets. A second tote bag contains 3 bottles of aspirin, 2 bottles of thyroid tablets and I bottle of laxative tablets. If 1 bottle of tablets is taken at random from each piece of luggage, find the probability that:

- (a) both bottles contain thyroid tablets: P(T, 1) T2)
- (b) neither bottle contains thyroid tablets; P(T, UT2)
- (c) the 2 bottles contain different tablets. = RITOU PUZ

A1: aspirin tablets are selected from the overnight case,

A2: aspirin tablets are selected from the tote bag,

L2: laxative tablets are selected from the tote bag,

T1: thyroid tablets are selected from the overnight case,

T2: thyroid tablets are selected from the tote bag.

(a) $P(T1 \cap T2) = P(T1)P(T2) = (3/5)(2/6) = 1/5$.

(b) $P(T'1 \cap T'2) = P(T'1)P(T'2) = (2/5)(4/6) = 4/15$.

(c) $1-P(A1 \cap A2)-P(T1 \cap T2) = 1-P(A1)P(A2)-P(T1)P(T2)$ =1-(2/5)(3/6)-(3/5)(2/6)=3/5.

71


Example 2.39: An electrical system consists of four components as illustrated in Figure 2.9. The system works if components A and B work and either of the components C or Dworks. The reliability (probability of working) of each component is also shown in Figure 2.9. Find the probability that (a) the entire system works and (b) the component C does not work, given that the entire system works. Assume that the four components work independently.

Solution: In this configuration of the system, A, B, and the subsystem C and D constitute a serial circuit system, whereas the subsystem C and D itself is a parallel circuit system.

(a) Clearly the probability that the entire system works can be calculated as $P[A \cap B \cap (C \cup D)] = P(A)P(B)P(C \cup D) = P(A)P(B)[1 - P(C' \cap D')]$ = P(A)P(B)[1 - P(C')P(D')]= (0.9)(0.9)[1 - (1 - 0.8)(1 - 0.8)] = 0.7776.

The equalities above hold because of the independence among the four com-

(b) To calculate the conditional probability in this case, notice that




Ex. 2.99

A circuit system is given in Figure 2.11. Assume the components fail independently.

(a) What is the probability that the entire system works?

(b) Given that the system works, what is the probability that the component A is not working?



Answer Ex. 2.99

(a)
$$P = 1 - [1 - (0.7)(0.7)][1 - (0.8)(0.8)(0.8)] = 0.75112.$$


P= 0.75112

$$(b) =$$

P= P(A'nCnDnE)

 $P_{\text{system works}}$ = (0.3)(0.8)(0.8)(0.8) = 0.2045 0.75112

 2.98 Suppose the diagram of an electrical system is given in Figure 2.10. What is the probability that the system works? Assume the components fail independently.



if
$$\Rightarrow P(B/S) = P(ANB'NCNPP) = 0.140840.95
system

given than

P(S)

P(S)$$

• EX:2.98

$$P = (0.95)[1 - (1 - 0.7)(1 - 0.8)](0.9) = 0.8037.$$

EX. 2.93

A town has 2 fire engines operating independently. The probability that a specific engine is available when needed is 0.96.

- (a) What is the probability that neither is available when needed? $P(A' \cap B) = 0.04*0.04=0.0016$
- (b) What is the probability that a fire engine is available when needed?

(a)
$$P(A' \cap B') = P(A')P(B') = (0.04)(0.04) = 0.0016.$$

(b) $P(A \cup B) = 1 - P(A' \cap B') = 1 - 0.0016 = 0.9984.$
(b) $P(A \cup B) = P(A') + P(B') - P(A \cap B)$

EX. 2.89 The probability that a doctor correctly diagnoses a particular illness is 0.7. Given that the doctor makes an incorrect diagnosis, the probability that the patient enters a law suit is 0.9. What is the probability that the doctor makes an incorrect diagnosis and the patient P(L)=0-9 sues?

$$8(2/5) = 0.9$$

 $8(500) = 8(210) = 9(500)$
 $9(500) = 0.340.9 = 0.27$

79

Answer of EX. 2.89:

 $P(A' \cap B) = P(A')P(B \mid A') = (0.3)(0.9) = 0.27$

If the events
$$A_1, A_2, \dots, A_k$$
 are independent, then
$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \cdots P(A_k|A_1 \cap A_2 \cap \dots \cap A_{k-1}).$$

$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1)P(A_2) \cdots P(A_k).$$

81

Bayes' Rule

عليها سؤالين

Table 2.1: Categorization of the Adults in a Small Town

Employed Unemployed Total 36 of Employees and Male 460 40 500 12 of Unemployed are Female 140 260400 members of the Total 600 300 900 Rotary Club $P(A) = P[(E \cap A) \cup (E' \cap A)] = P(E \cap A) + P(E' \cap A)$ P(E)P(A|E) + P(E')P(A|E')Find the probability of the event A that the P(A) individual selected is a member of the Rotary Club? ENA A= Ena UE' DA

Figure 2.12: Venn diagram for the events A, E, and E'.

				3
NEW .	800 - T	P(A(E)	000	

	100.00			
				1
24.8		A(E') =	300	
			_	

. 1	P(AE) = 350	PROPIAGO
23/		
Z		
5/		* P(E)P(A'E)

ble 2.5	- Lance	Unemployed	Total
	Employed	40	500
Make	460	260	400
Founde	140	300	900
Total	0.00		

36 of Employees and 12 of Unemployed are members of the **Rotary Club**

Figure 2.13: Tree diagram for the data on page 63, using additional information on page 72.

the probability P(E')P(A|E'), it follows that

$$P(A) = \left(\frac{2}{3}\right)\left(\frac{3}{50}\right) + \left(\frac{1}{3}\right)\left(\frac{1}{25}\right) = \frac{4}{75}.$$

83

Theorem 2.13: If the events B_1, B_2, \dots, B_k constitute a partition of the sample space S such that $P(B_i) \neq 0$ for $i = 1, 2, \dots, k$, then for any event A of S,

$$P(A) = \sum_{i=1}^k P(B_i \cap A) = \sum_{i=1}^k P(B_i) P(A|B_i).$$




Figure 2.14: Partitioning the sample space S.

The event A is seen to be the union of the mutually exclusive events

Proof: Consider the Venn diagram of Figure 2.14. The event A is seen to be the union of the mutually exclusive events

$$B_1 \cap A$$
, $B_2 \cap A$, ..., $B_k \cap A$;

that is,

$$A = (B_1 \cap A) \cup (B_2 \cap A) \cup \cdots \cup (B_k \cap A).$$

Using Corollary 2.2 of Theorem 2.7 and Theorem 2.10, we have

$$P(A) = P[(B_1 \cap A) \cup (B_2 \cap A) \cup \dots \cup (B_k \cap A)]$$

$$= P(B_1 \cap A) + P(B_2 \cap A) + \dots + P(B_k \cap A)$$

$$= \sum_{i=1}^k P(B_i \cap A)$$

$$= \sum_{i=1}^k P(B_i) P(A|B_i).$$

85

Example 2.41: In a certain assembly plant, three machines, B_1 , B_2 , and B_3 , make 30%, 45%, and 25%, respectively, of the products. It is known from past experience that 2%, 3%, and 2% of the products made by each machine, respectively, are defective. Now, suppose that a finished product is randomly selected. What is the probability that it is defective?

Solution: Consider the following events:

A: the product is defective,


 B_1 : the product is made by machine B_1 ,

 B_2 : the product is made by machine B_2 ,

 B_3 : the product is made by machine B_3 .


Applying the rule of elimination, we can write

$$P(A) = P(B_1)P(A|B_1) + P(B_2)P(A|B_2) + P(B_3)P(A|B_3).$$



B,=	0	M3
BZ	=	0.45
		0.25

A



Referring to the tree diagram of Figure 2.15, we find that the three branches give the probabilities

$$P(B_1)P(A|B_1) = (0.3)(0.02) = 0.006,$$

 $P(B_2)P(A|B_2) = (0.45)(0.03) = 0.0135,$
 $P(B_3)P(A|B_3) = (0.25)(0.02) = 0.005,$

and hence

$$P(A) = 0.006 + 0.0135 + 0.005 = 0.0245$$
.

P(P) = P(D,) +P(P2) +P(200 as less 1 200) 200 as less 1 200) 200 as less 1 2000 2 0.006+0.0135+0 = 0.0245

عدم الأل المحداد لوخرال (B/D)

Theorem 2.14: (Bayes' Rule) If the events B_1, B_2, \dots, B_k constitute a partition of the sample space S such that $P(B_i) \neq 0$ for $i = 1, 2, \dots, k$, then for any event A in S such that $P(A) \neq 0$

$$P(A) \neq 0$$
,
 $P(B_r|A) = \frac{P(B_r \cap A)}{\sum_{i=1}^{k} P(B_i \cap A)} = \frac{P(B_r)P(A|B_r)}{\sum_{i=1}^{k} P(B_i)P(A|B_i)}$ for $r = 1, 2, ..., k$.

Proof: By the definition of conditional probability.

$$P(B_r|A) = \frac{P(B_r \cap A)}{P(A)}$$

and then using Theorem 2.13 in the denominator, we have
$$P(B_r|A) = \frac{P(B_r \cap A)}{\sum\limits_{i=1}^k P(B_i \cap A)} = \frac{P(B_r)P(A|B_r)}{\sum\limits_{i=1}^k P(B_i)P(A|B_i)}$$

which completes the pro-

Example 2.42: With reference to Example 2.41, if a product was chosen randomly and found to be defective, what is the probability that it was made by machine B_3 ?

Solution: Using Bayes' rule to write

$$P(B_3|A) = \frac{P(B_3)P(A|B_3)}{P(B_1)P(A|B_1) + P(B_2)P(A|B_2) + P(B_3)P(A|B_3)}$$

and then substituting the probabilities calculated in Example 2.41, we have

bestituting the probabilities calculated in Example 2.005
$$P(B_3|A) = \frac{0.005}{0.006 + 0.0135 + 0.005} = \frac{0.005}{0.0245} = \frac{10}{49}$$
. So that

In view of the fact that a defective product was selected, this result suggests that it probably was not made by machine B_3 .

87

Example 2.43: A manufacturing firm employs three analytical plans for the design and development of a particular product. For cost reasons, all three are used at varying times. In fact, plans 1, 2, and 3 are used for 30%, 20%, and 50% of the products, respectively. The defect rate is different for the three procedures as follows:

$$P(D|P_1) = 0.01$$
, $P(D|P_2) = 0.03$, $P(D|P_3) = 0.02$,

where $P(D|P_1) = 0.01$, $P(D|P_2) = 0.03$, $P(D|P_3) = 0.02$, $P(D|P_3) = 0.02$, where $P(D|P_3)$ is the probability of a defective product, given plan j. If a random product was observed and found to be defective, which plan was most likely used and thus responsible?

Solution: From the statement of the problem P(D|D)Solution: From the statement of the problem P (P/D)

$$P(P_1) = 0.30$$
, $P(P_2) = 0.20$, and $P(P_3) = 0.50$,

we must find $P(P_j|D)$ for j = 1, 2, 3. Bayes' rule (Theorem 2.14) shows

$$P(P_1|D) = \frac{P(P_1)P(D|P_1)}{P(P_1)P(D|P_1) + P(P_2)P(D|P_2) + P(P_3)P(D|P_3)}$$


$$= \frac{(0.30)(0.01)}{(0.3)(0.01) + (0.20)(0.03) + (0.50)(0.02)} = \frac{0.003}{0.019} = 0.158. \text{ M}$$


Similarly.

$$P(P_2|D) = \frac{(0.03)(0.20)}{0.019} = 0.316 \text{ and } P(P_3|D) = \frac{(0.02)(0.50)}{0.019} = 0.526.$$

The conditional probability of a defect given plan 3 is the largest of the three; thus a defective for a random product is most likely the result of the use of plan 3.


Using Bayes' rule, a statistical methodology called the Bayesian approach has attracted a lot of attention in applications. An introduction to the Bayesian method will be discussed in Chapter 18.





Definitions:

- The lower half of a data set is the set of all values that are to the left of the median value when the data has been put into increasing order.
- The upper half of a data set is the set of all values that are to the right of the median value when the data has been put into increasing order.
- The first quartile, denoted by Q_1 , is the median of the lower half of the data set. This means that about 25% of the numbers in the data set lie below Q_1 and about 75% lie above Q_1 .
- The third quartile, denoted by Q_3 , is the median of the upper half of the data set. This means that about 75% of the numbers in the data set lie below Q_3 and about 25% lie above Q_3 .



Le 15

تنعق عندال 1.5

IPR-+36

لعين

25 (n+1)+025 Location value (CBO of Find the first and third quartiles of the data set {3, 7, 8, 5, 12, 14, 21, 13, 16, 18}. in increasing order: 3, 5, 7, 8, 12, 13, 14, 16, 18, 21. Location of Q1: (10+1)*0.25=2.75 n4 Interpolation Q1=value of location 2+0.75*(value of location 3- value of location 2) Q1=5+0.75*(7-5)=6.5 Location of Q2: (10+1)*0.5=5.5 Q2= (12+13)/2=12.5 Location of Q3: (10+1)*0.75=8.25 25 10361 Q3=value of location 8+0.25*(value of location 9- value of location 8) Q3=16+0.25*(18-16)=16.5 Inter quartile range (IQR)= Q3-Q1 9.3-9.1 Q 3 UL= 16.5-6.5=10 > 7 9 K UL= 16.5+ (1.5*10) = 31.5 > Q3 > L(n+1)+0.75 1 11+0.75=8.25 LL=6.5-(1.5*10)=31.5 Q. 1 | 1.5*10) = -8.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1.5 | 1. point

4) Box-and-Whisker Plot or Box Plot -You have to know to estimate the percentile and quartile

e.g., Nicotine content was measured in a random sample of 40 cigarettes. The data are displayed in Table 1.8.

Table 1.8: Nicotine Data for Example 1.5

		0.01	1.70	2.28	1.74	1.47	1.97
			1.79	1 70	2.17	2.55	2.11
0.85	1.24						1.85
1.86	1.90	1.68	1.51		0.72		1.93
1.82	1.79	2.46	1.88	2.08	1.67	1.37	1.00
1.40	1.64	2.09	1.75	1.63	2.37	1.75	1.69

In order

lolder								4	of ha
-	702	3	81	5	5	7	1 51	1 58	1.63 1.75
0.72	0.85	1.09	1.24	1.37	1.4	1.4/	1.51	1.50	175-
11160	164	1.67	1.60	1 69	1 69	1.7	1.74	1.75	1.75
1.04	121.04	1.07	1.00	1.05	1.00	10	1 02	1 93	1.97 30
2 1.79	1.79	1.82	1.85	1.86	1.88	1.9	1.52	1.50	1.97 3°°
2.03	2.08	2.09	2.11	2.17	2.28	2.31	2.37	2.46	2.55
31									





Figure 1.9: Box-and-whisker plot for Example 1.5.



12) 3 11 jul & 11 Jones

• Find 62% percentile of {3, 7, 8, 5, 12, 14; 21, Location 3/5/7/8/12/13/14/16/18/21 13, 16, 18}.

Location of 62% percentile is (n+1)*0.62=6.82

62% percentile=Value of location 6+0.82*(Value of location 7-Value of location 6)=13+0.82*(14-13)=13.82

134082/14-18

Find 29% percentile of Table 1.8 n=40

Location of 29% percentile is (40+1)*0.29=11.89


29% percentile=Value of location 11+0.89*(Value of location 12-Value of location 11)=1.64+0.29*(1.64-1.64)=1.64

عن الراق المحاصر المرقة عن الرقة المرقة الم

1 PH+0.89 (6)

29

Slide 22 Jewick Digen also III frequency/12003/51 1) f = B 25 = 40 = 0.125 RA = 1-0.5-0.L. (U



اليان سليمان احمد ياسين



Probability and Statistics

به عمان احتمالیه حدل معین

Chapter 2

Department of Industrial Engineering

Sample singulary space singulary space singulary space singulary space singulary spaces spaces singulary spaces s

- Statisticians use the word experiment to describe any process that generate a set of data.
- Example of a statistical experiment is the tossing of a coin.

S ~ Sample Space

الفغاء العبني

aules Diasil element les Mis (so lipse assesso

The set of all possible outcomes of a statistical experiment is called the sample Definition 2.1: space and is represented by the symbol S.

> Each outcome in a sample space is called an element or a member of the sample space, or simply a sample point. If the sample space has a finite number of elements, we may list the members separated by commas and enclosed in braces. Thus, the sample space S, of possible outcomes when a coin is inpped, may be written

COIN = S (H.T).

where H and T correspond to heads and tails, respectively.

Lesling, reisi N Estrellaviol cello iso lo 50 doi la 10 10 - M - W - WIED 11

1

Example 2.1: Consider the experiment of tossing a die. If we are interested in the number that shows on the top face, the sample space is

 $S_1 = \{1, 2, 3, 4, 5, 6\}.$

If we are interested only in whether the number is even or odd, the sample space is simply

النزد (even, odd). حر النزد

** More than one sample space can be used to describe the outcomes of an experiment.

In some experiment it is helpful to list the elements of the sample space by means of a tree diagram.

> Grappe Jietz Ust bischousen space complerated

4

التجاز بالإحمانية نسطيع التمسر عنفا بأغرض طريقه مله ١١ كالحر النر رص ١-١

Example 2.2 An experiment consists of flipping a coin and then flipping it a second time if a head occurs. If a tail occurs on the first flip, then a die is tossed once. To list the elements of the sample space providing the most information, we construct the tree diagram of Figure 2.1. The various paths along the branches of the tree give the distinct sample points. Starting with the top left branch and moving to the right along the first path, we get the sample point HH, indicating the possibility that heads occurs on two successive flips of the coin. Likewise, the sample point T3 indicates the possibility that the coin will show a tail followed by a 3 on the toss of the die. By proceeding along all paths, we see that the sample space is

 $S = \{HH, HT, T1, T2, T3, T4, T5, T6\}.$

dipgram

dipgram

complecated sample space estable

coin




Figure 2.1: Tree diagram for Example 2.2.

5

Propability: Unison solyphaste Sample space

المعنى عسورات مساورة عورات مساورة المعاددة المع

Example 2.3: Suppose that three items are selected at random from a manufacturing process. Each item is inspected and classified defective, <u>D</u>, or <u>nondefective</u>, <u>N</u>. To list the elements of the sample space providing the most information, we construct the tree diagram of Figure 2.2. Now, the various paths along the branches of the tree give the distinct sample points. Starting with the first path, we get the sample point <u>DDD</u>, indicating the possibility that all three items inspected are defective. As we proceed along the other paths, we see that the sample space is

 $S = \{DDD, DDN, DND, DNN, NDD, NDN, NND, NNN\}.$

for too items




Figure 2.2: Tree diagram for Example 2.3.

(N) 8 mol

فالدة ال ماه الخالف المونى ال

A rule plient element Telis ice évalement

• Sample spaces with a large or infinite number of sample points are best described by a statement or rule.

• For example, if the possible outcomes of an experiment are the set of cities in the world with

a. population over 1 million:

 $S = \{x \mid x \text{ is a city with a. population over 1 million}\},$ which reads "S is the set of all x such that x is a city with a population over 1 million.

Us a city with al

Sample 1105/2000 Events

Definition 2.2: An event is a subset of a sample space.

 $S = \{DDD, DDN, DND, DNN, NDD, NDN, NND, NNN\}.$

• Event B represents the number defectives is greater than 1 ->

 $= \{DDN, DND, NDD, DDD\}$ $S = \{1/2/3/4/5/6\}$ like more than

at less one > villes 3 de 7 = A = 12,1, vient lemes 210 8

at least = = 3DNN/NDN/NND/NNNS two N

والمكار و المحار المحار و ال

electronic component, then the event A that the component fails before the end of the fifth year is the subset $A = \{t \mid 0 \le t < 5\}$. Dolalmous S 200161

Definition 2.3: The complement of an event A with respect to S is the subset of all elements of S that are not in A. We denote the complement of A by the symbol A'.

Example 2.6: Consider the sample space

 $S = \{\text{book, cell phone, mp3, paper, stationery, laptop}\}.$

Let $A = \{book, stationery, laptop, paper\}$. Then the complement of A is A' ={cell phone, mp3}.

S={ 1,2,3,4,5,6}


If event A is the even number in tossing of a die

 $A=\{2,4,6\}, A'=\{1,3,5\}$

more than If event B is a number greater than 3 in a tossing of a die

 $B=\{4,5,6\}, B'=\{1,2,3\}$ 3 (10 25 i

review (1003) 3)12




Example 2.12: If $M = \{x \mid 3 < x < 9\}$ and $N = \{y \mid 5 < y < 12\}$, then $M \cup N = \{z \mid 3 < z < 12\}. \qquad M \cap N = \{z \mid 5 \mid 5 \mid 12\}$ He disdoint event \rightarrow continuous (selecute to featral What

The person of the continuous (selecute to featral What

The person of the continuous (selecute to featral What \rightarrow Michigan (the leant) \rightarrow Michigan \rightarrow

 The relationship between events and the corresponding sample space can be illustrated graphically by means of Venn diagrams.



13


The relationship between events and the corresponding sample space can be illustrated graphically by means of Venn diagrams. In a Venn diagram we let the sample space be a rectangle and represent events by circles drawn inside the rectangle. Thus, in Figure 2.3, we see that

$$A \cap B = \text{regions 1 and 2},$$

 $B \cap C = \text{regions 1 and 3},$

$$A = \begin{cases} 1/2/4/7 \\ B = \begin{cases} 1/2/3/63 \end{cases}$$

$$C = \begin{cases} 1/3/4/63 \end{cases}$$



event 11 Mai jul 1

Figure 2.3: Events represented by various regions.

$$\begin{cases}
A \cup C = \text{ regions } 1, 2, 3, 4, 5, \text{ and } 7, \\
A \cap B \cap C = \frac{\text{region } 1}{\text{regions } 2, 6, \text{ and } 7}, \\
A \cup B \cap C' = \frac{\text{region } 2, 6}{\text{regions } 2, 6, \text{ and } 7},
\end{cases}$$

$$\begin{cases}
112141713163 \cap \left\{21716\right\}
\end{cases}$$

$$= \left\{21617\right\}$$




Figure 2.4: Events of the sample space S.

In Figure 2.4, we see that events A, B, and C are all subsets of the sample space S. It is also clear that event B is a subset of event A; event $B \cap C$ has no elements and hence B and C are mutually exclusive; event $A \cap C$ has at least one element; and event $A \cup B = A$. Figure 2.4 might, therefore, depict a situation where we select a card at random from an ordinary deck of 52 playing cards and observe whether the following events occur:


A: the card is red.


B: the card is the jack, queen, or king of diamonds,

C: the card is an ace.

Clearly, the event $A \cap C$ consists of only the two red aces.

Several results that follow from the foregoing definitions, which may easily be verified by means of Venn diagrams, are as follows:





15

• Exercise 2.7:

Four students are selected at random from a chemistry class and classified as male or female. List the elements of the sample space S_1 using the letter M for "male" and F for "female." Define a second sample space S_2 where the elements represent the number of females selected.

S₁={MMMM,MMMF,MMFM,MFMM, FMMM,MMFF,MFMF,MFFM,FMFM,FFMM, FMMF,MFFF,FMFF,FFMF,FFFM,FFFF}

 $S_2 = \{0, 1, 2, 3, 4\}$ Sample space

Sample space

Sample space

Sample space

Ship for all 1923

Semale 1923

tree free

17

• Exercise 2.14:

If $S = \{0,1,2,3,4,5,6,7,8,9\}$ and $A = \{0,2,4,6,8\}$, $B = \{1,3,5,7,9\}$, $C = \{2,3,4,5\}$, and

$$D = \{1, 6, 7\}$$

- (a) $A \cup C := \{0,2,3,4,5,6,8\}.$
- (b) $A \cap B = \phi$.
- (c) C'; = {0,1,6,7,8,9}.
- (d) $(C' \cap D) \cup B$; $C' \cap D = \{1,6,7\}$, $(C' \cap D) \cup B = \{1,3,5,6,7,9\}$
- (e) $(S \cap C)'$; $C' = \{0,1,6,7,8,9\}$. $(5 \cup C)$
- (f) AnCnD': {2,4}, so AnC nD' = {2,4}.

Rule 2.1: If an operation can be performed in n_1 ways, and if for each of these ways a second operation can be performed in n_2 ways, then the two operations can be performed together in n_1n_2 ways,

thrown once?

عنارات عاملة = خيارات به حيالة عاملة عامل


Solution: The first die can land face-up in any one of $n_1 = 6$ ways. For each of these 6 ways, the second die can also land face-up in $n_2 = 6$ ways. Therefore, the pair of dice can land in $n_1 n_2 = (6)(6) = 36$ possible ways.

Example 2.14: A developer of a new subdivision offers prospective home buyers a choice of Tudor, rustic, colonial, and traditional exterior styling in ranch, two-story, and split-level floor plans. In how many different ways can a buyer order one of these homes?

Solution:

Since $n_1 = 4$ and $n_2 = 3$, a buyer must choose from

 $n_1 n_2 = (4)(3) = 12$ possible homes.






Figure 2.6: Tree diagram for Example 2.14.

Q.

Example 2.15: If a 22-member club needs to elect a chair and a treasurer, how many different ways can these two to be elected?

22 Solution Chair & treasure

Solution: For the chair position, there are 22 total possibilities. For each of those 22 possibilities, there are 21 possibilities to elect the treasurer. Using the multiplication rule, we obtain $n_1 \times n_2 = 22 \times 21 = 462$ different ways.

The multiplication rule, Rule 2.1 may be extended to cover any number of operations. Suppose, for instance, that a customer wishes to buy a new cell phone and can choose from $n_1 = 5$ brands, $n_2 = 5$ sets of capability, and $n_3 = 4$ colors. These three classifications result in $n_1n_2n_3 = (5)(5)(4) = 100$ different ways for a customer to order one of these phones. The generalized multiplication rule covering k operations is stated in the following.

Rule 2.2:

If an operation can be performed in n_1 ways, and if for each of these a second operation can be performed in n_2 ways, and for each of the first two a third operation can be performed in n_3 ways, and so forth, then the sequence of k operations can be performed in $n_1 n_2 \cdots n_k$ ways.

Example 2.16: Sam is going to assemble a computer by himself. He has the choice of chips from two brands, a hard drive from four, memory from three, and an accessory bundle from five local stores. How many different ways can Sam order the parts?

Solution: Since $n_1 = 2$, $n_2 = 4$, $n_3 = 3$, and $n_4 = 5$, there are

 $n_1 \times n_2 \times n_3 \times n_4 = 2 \times 4 \times 3 \times 5 = 120$

different ways to order the parts.


5×5×4 = 100 U)

brand H.D menocal

2 4 4 4 3 45

= 120

2.22 In a medical study patients are classified in 8 ways according to whether they have blood type AB^+ , AB^- , A^- , A^- , B^+ , B^- , O^+ , or O^- , and also according to whether their blood pressure is low, normal, or high. Find the number of ways in which a patient can be classified.




21

Answer (Ex. 22)

n₁n₂= 8*3=24 classification

2.27 A developer of a new subdivision offers a prospective home buyer a choice of 4 designs. 3 different heating systems, a garage or carport, and a patio or screened porch. How many different plans are available to this buyer?



23

Answer (Ex. 27)

 $n_1 n_2 n_3 n_4 = 4*3*2*2=48$ different house plans

20 La Colo Caroli

lice that the license number contained the letters RLH ness cannot recall the last 2 digits but is certain that all 3 digits are different find the maximum number of check.

RLH 5[--]
5 Cho 47

6 Cho

(3) En lot 151 x - - 9 8


Caro 5 5 1 1 Kess 10 9 8

Similar

Answer (Ex. 33)

 $n_1n_2 = 9*8=72$ registrations to be checked.

Example 4



(a) There are no restrictions?

(b) A will serve only if he is president?

(c) B and C will serve together or not at all?

Solution:

(a)
$$_{50}P_2 = \frac{501}{(50-2)!} = \frac{50!}{48!} = \frac{50 \cdot 49 \cdot 48!}{48!} = 50 \cdot 49 = 2450$$

- (b) Since A will serve only if he is president, we have two situations here: A is selected as the president, which yields 49 possible outcomes for the treasurer's position, or officers are selected from the remaining 49 people without A, which has the number of choices $_{49}P_2 = 49 \times 48 = 2352$. Therefore, the total number of choices is 49 + 2352 = 2401.
- (c) The number of selections when B and C serve together is 2. The number of selections when both B and C are not chosen is $_{48}P_2 = 2256$. Therefore, the total number of choices in this situation is 2 + 2256 = 2258.

Example

- A president and a treasurer are to be chosen from a student club consisting of 50 people.
 How many different choices of officers are possible if
 - (c) B and C will serve together or not at all:

(1) B present.
$$C - Treasmen - 1$$
 $C = Presid$
 $N = 48$, $N = 2$
 $N = 48$, $N = 2$

27

Example:

How many even three digit numbers can be performed from the digits 1,2,5,6 and 9 if each digit can be used only once?

$$n_1 n_2 n_3 = (4)(3)(2) = 24$$

(معنى ترى ع) منطقة الزال 0/1/2/5/6/9 mple 2.17: How many even four-digit numbers can be formed from the digits 0, 1, 2, 5, 6, and 9 if each digit can be used only once? **Solution:** Since the number must be even, we have only $n_1 = 3$ choices for the units position. However, for a four-digit number the thousands position cannot be 0. Hence, we consider the units position in two parts, 0 or not 0. If the units position is 0 (i.e., $n_1 = 1$), we have $n_2 = 5$ choices for the thousands position, $n_3 = 4$ for the hundreds position, and $n_4 = 3$ for the tens position. Therefore, in this case we have a total $n_1 n_2 n_3 n_4 = (1)(5)(4)(3) = 60$ even four-digit numbers. On the other hand, if the units position is not 0 (i.e., $n_1 = 2$), we have $n_2 = 4$ choices for the thousands position, $n_3 = 4$ for the hundreds position, and $n_4 = 3$ for the tens position. In this situation, there are a total of $n_1 n_2 n_3 n_4 = (2)(4)(4)(3) = 96$ Since the above two cases are mutually exclusive of each other, the total number 0/1 of even four-digit numbers can be calculated by 60 + 96 = 156Definition 2.7: A permutation is an arrangement of all or part of a set of objects. Consider the three letters a, b, and c. The possible permutations are abc, acb, bac, bca, cab, and cba. Thus, we see that there are 6 distinct arrangements. Using Rule 2.2, we could arrive at the answer 6 without actually listing the different orders by the following arguments: There are $n_1 = 3$ choices for the first position. No matter which letter is chosen, there are always $n_2 = 2$ choices for the second position. No matter which two letters are chosen for the first two positions, there is only $n_3 = 1$ choice for the last position, giving a total of $n_1 n_2 n_3 = (3)(2)(1) = 6$ permutations by Rule 2.2. In general, n distinct objects can be arranged in linear $n(n-1)(n-2)\cdots(3)(2)(1)$ ways. There is a notation for such a number. 1) no biect states n! - so abc 3) nobuect bis different object **Theorem 2.1:** The number of permutations of n objects is n!. The number of permutations of the four letters a, b, c, and d will be 4! = 24. Now consider the number of permutations that are possible by taking two letters at a time from four. These would be ab, ac, ad, ba, bc, bd, ca, cb, cd, da, db, and

dc. Using Theorem 2.1 again, we have two positions to fill with $n_1 = 4$ choices for the first and then $n_2 = 3$ choices for the second for a total of

$$n_1n_2 = (4)(3) = 12$$

permutations. In general, n distinct objects taken r at a time can be arranged in

$$n(n-1)(n-2)\cdots(n-r+1)$$


ways. We represent this product by the symbol

$$_{n}P_{r}^{s}=\frac{n!}{(n-r)!}.$$

As a result we have the theorem that follows.

The number of permutations of n distinct objects taken r at a time is Theorem 2.2:

$$\int_{n} P_r = \frac{n!}{(n-r)!}.$$



واجن کی فی اکل

Example 2.18: In one year, three awards (research, teaching, and service) will be given to a class of 25 graduate students in a statistics department. If each student can receive at most one award, how many possible selections are there?

Solution: Since the awards are distinguishable, it is a permutation problem. The total number of sample points is

$$N = 25$$

$$R = \frac{25!}{(25-3)!} = \frac{25!}{22!} = (25)(24)(23) = 13,800.$$

$$25 = \frac{25!}{25!} = \frac{25 \times 24 \times 23!}{23!}$$

$$= 13800$$

31

a bleo 9

+660001

Example 2.19: A president and a treasurer are to be chosen from a student club consisting of 50 people. How many different choices of officers are possible if

(a) there are no restrictions;

(b) A will serve only if he is president; >> projection by

(d) D and E will not serve together?

Solution: (a) The total number of choices of officers, without any restrictions, is

$$_{50}P_2 = \frac{50!}{48!} = (50)(49) = 2450.$$

(b) Since A will serve only if he is president, we have two situations here: (i) A is selected as the president, which yields 49 possible outcomes for the treasurer's position, or (ii) officers are selected from the remaining 49 people without A, p which has the number of choices $_{49}P_2 = (49)(48) = 2352$. Therefore, the total number of choices is $_{49}P_2 = 2401$.

(c) The number of selections when B and C serve together is 2. The number of selections when both B and C are not chosen is $_{48}P_2 = 2256$. Therefore, the total number of choices in this situation is 2 + 2256 = 2258.

(d) The number of selections when D serves as an officer but not E is (2)(48) = 96, where 2 is the number of positions D can take and 48 is the number of selections of the other officer from the remaining people in the club except E. The number of selections when E serves as an officer but not D is also (2)(48) = 96. The number of selections when both D and E are not chosen is ${}_{48}P_2 = 2256$. Therefore, the total number of choices is (2)(96) + 2256 = 2148. This problem also has another short solution: Since D and E can only serve together in 2 ways, the answer is 2450 - 2 = 2448.

101/2 = 16 = 24 L

Start again with the 2450 permutations. If B 1

other than C who might be treasurer. And if E other than C who might be president. We multiple also eliminate the 96 outcomes in which C w 2450 - 96 - 96 = 2258

pt

so
en

other than C who might be treasurer. And if E other than C who might be president. We multiple also eliminate the 96 outcomes in which C w 2450 - 96 - 96 = 2258

The number of selections when both D and E are not chosen. Therefore, the total number of choices is (2)(96) + 2256 = blem also has another short solution: Since D and E can only in 2 ways, the answer is 2450 - 2 = 2448.


22 48×47 11 + 1=2

الصدوهمال كا

2 48+47

Theorem 2.3: The number of permutations of n objects arranged in a circle is (n-1)!.

So far we have considered permutations of distinct objects. That is, all the objects were completely different or distinguishable. Obviously, if the letters b and objects were completely different or distinguishable. Obviously, if the letters b and c become c are both equal to x, then the 6 permutations of the letters a, b, and c become axx, axx, xax, xax, xxa, and xxa, of which only 3 are distinct. Therefore, with 3 axx, axx, xax, xax, xax, xax, xax, xax, xax, and xxa, of which only 3 are distinct. Therefore, with 3 axx, axx




35

Example 2.20: In a college football training session, the defensive coordinator needs to have 10 players standing in a row. Among these 10 players, there are 1 freshman, 2 sophomores, 4 juniors, and 3 seniors. How many different ways can they be arranged in a row if only their class level will be distinguished?

Solution: Directly using Theorem 2.4, we find that the total number of arrangements is

$$\frac{10!}{1! \ 2! \ 4! \ 3!} = 12,600.$$



Theorem 2.5: The number of ways of partitioning a set of n objects into r cells with n_1 elements in the first cell, n_2 elements in the second, and so forth, is

$$\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!},$$

where $n_1 + n_2 + \dots + n_r = n$.

Example 2.21: In how many ways can 7 graduate students be assigned to 1 triple and 2 double hotel rooms during a conference?

Solution: The total number of possible partitions would be

$$\binom{7}{3,2,2} = \frac{7!}{3!\ 2!\ 2!} = 210.$$

Juis Mosto

Example 2.23: How many different letter arrangements can be made from the letters in the word

Solution: Using the same argument as in the discussion for Theorem 2.6, in this example we can actually apply Theorem 2.5 to obtain

$$\binom{10}{3,3,2,1,1} = \frac{10!}{3!\ 3!\ 2!\ 1!\ 1!} = 50,400.$$

Here we have 10 total letters, with 2 letters (S, T) appearing 3 times each, letter I appearing twice, and letters A and C appearing once each. On the other hand, this result can be directly obtained by using Theorem 2.4.

37

In many problems, we are interested in the number of ways of selecting r objects from n without regard to order. These selections are called **combinations**. A combination is actually a partition with two cells, the one cell containing the r objects selected and the other cell containing the (n-r) objects that are left. The number of such combinations, denoted by

$$\binom{n}{r,n-r}$$
, is usually shortened to $\binom{n}{r}$,

since the number of elements in the second cell must be n-r.

The number of combinations of n distinct objects taken r at a time is Theorem 2.6:

select $\binom{n}{r} = \frac{n!}{r!(n-r)!}$

Example 2.22: A young boy asks his mother to get 5 Game-Boy TM cartridges from his collection of 10 arcade and 5 sports games. How many ways are there that his mother can get 3 arcade and 2 sports games?

Solution: The number of ways of selecting 3 cartridges from 10 is


 $\binom{10}{3} = \frac{10!}{3! (10-3)!} = 120.$

The number of ways of selecting 2 cartridges from 5 is

$$\binom{5}{2} = \frac{5!}{2! \ 3!} = 10.$$

Using the multiplication rule (Rule 2.1) with $n_1 = 120$ and $n_2 = 10$, we have

Osing the multiplication rule (Rule 2.1) with $n_1 = 120$ and $n_2 = 10$, where $n_2 = 120$ and $n_3 = 120$ and $n_4 = 120$ and $n_$



Permutations and Combinations

Number of permutations (order matters) of *n* things taken *r* at a time:

$$P(n,r) = \frac{n!}{(n-r)!}$$

Number of combinations (order does not matter) of *n* things taken *r* at a time:

$$C(n,r) = \frac{n!}{(n-r)!r!}$$

Number of different permutations of n objects where there are n_1 repeated items, n_2 repeated items, ... n_k repeated items

$$\frac{n!}{n_1!n_2!...n_k!}$$

• If an experiment consists of throwing a die and then drawing a letter at random from the English alphabet, how many points are there in the sample space?

41

Ex 2.32:

a) How many distinct permutations can be made from the letters of the word columns?,

5040

b) How many of these permutations start with the letter m:

columns (3 m 620)

6!=720

ab cd * 4

EX. 2.31

If a multiple-choice test consists of 5 questions each with 4 possible answers of which only 1 is correct,

(a) In how many different ways can a student check off one answer to each question?

(45) = 1024

(b) In how many ways can a student, check off one answer to each question and get all the answers wrong?


43

Ex. 2.40

In how many ways can 5 starting positions on a basketball team be filled with 8 men who can play any of the positions?

1.1 Overview: Statistical Inference, Samples, Populations, and Experimental Design

- -Use of Scientific Data
- -Variability in Scientific Data
- -The Role of Probability



إ مِلْ سَرَرِ حِينَ لَسَجُرِ لِللَّهِ عِلَى

Example 1.2

3

Often the nature of the scientific study will dictate the role that probability and deductive reasoning play in statistical inference. Exercise 9.40 on page 297 provides data associated with a study conducted at the Virginia Polytechnic Institute and State University on the development, of a relationship between the roots of trees and the action of a fungus. Minerals are transferred from the fungus to the trees and sugars from the trees to the fungus. Two samples of 10 northern red oak seedlings are planted in a greenhouse, one containing seedlings treated with nitrogen and one containing no nitrogen. All other environmental conditions are held constant. All seedlings contain the fungus *Pisolithus tinctorus*. More details are supplied in Chapter 9. The stem weights in grams were recorded after the end of 140 days. The data are given in Table 1.1.


المنجرة

و درسانالولا ۱ urpose of this experiment: the use of nitrogen has an influence on the growth of the roots. Table 1.1: Data Set for Example 1.2 No Nitrogen برزه البالات نعراله را بهافته الم Nitrogen 0.32 0.26 - 51 ds1-0.53 0.43 Jo 6 0.28 0.47 0.37 0.49 0.52 (0.26-0.86) 0.47 e 15) 14/4/2 cm? 0.43 0.75 0.36 0.79 0.86 - de i 0.42 (0.28-0.53) 0.38 0.62 0.43 0.46 0.35 0.40 0.45 0.50 0.55 0.65 0.75 0.60 0.70 0.80 1053 م دراس تا شر طدة و احدة Figure 1.1: A dot plot of stem weight data. بتثبيت بافي العوامل GAST Shows summer statistics 9 Je the Me

+ الإطاعة يجمع أن نكن مدروسة وليس عنوائية

- On average, the use of nitrogen increases the stem weight.
- Nitrogen is effective but ,how can this be quantified. ?

The 3 dol ishila of the Bird and is to



1.3 Sampling Procedures; Collection of Data faille mésus revels e 1000 cycle jes i 15 to 4 4000 Timor prisont cycle Timor prisont

Simple Random Sampling

Experimental Design

Example 1.3:1 A corrosion study was made in order to determine whether corrosion of an aluminum metal coated with a corrosion retardation substance reduced the amount of corrosion. The coating is a protectant that is advertised to minimize fatigue damage in this type of material. Also of interest is the influence of humidity on the amount of corrosion. A corrosion measurement can be expressed in thousands of cycles to failure. Two levels of coating, no coating and chemical corrosion coating, were used. In addition, the two relative humidity levels are 20% relative humidity and 80% relative humidity.

धंरण राजमां। (पर्वाक , त्वार) । विर्वाद

المع من عددالدولاات

Lew / Lero منحلال عرد المدران كل مار ١١عدد الدوران، قل

1.4 Measures of Location: The Sample Mean and Median

Definition 1.1: Suppose that the observations in a sample are x_1, x_2, \ldots, x_n . The sample mean, denoted by \bar{x} , is

$$\bar{x} = \sum_{i=1}^{n} \frac{x_i}{n} = \frac{x_1 + x_2 + \dots + x_n}{n}.$$

Definition 1.2: Given that the observations in a sample are x_1, x_2, \ldots, x_n , arranged in increasing order of magnitude, the sample median is

$$\tilde{x} = \begin{cases} x_{(n+1)/2}, & \text{if } n \text{ is odd,} \\ \frac{1}{2}(x_{n/2} + x_{n/2+1}), & \text{if } n \text{ is even.} \end{cases}$$

11

As an example, suppose the data set is the following: 1.7, 2.2, 3.9, 3.11, and 14.7. The sample mean and median are, respectively,

The sample mean and median are, respectively,
$$\bar{x} = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

$$|\hat{x}| = 5.12, \quad \hat{x} = 3.9. \quad 3.11$$

Clearly, the mean is influenced considerably by the presence of the extreme observation, 14.7, whereas the median places emphasis on the true "center" of the data set. In the case of the two-sample data set of Example 1.2, the two measures of central tendency for the individual samples are

$$\bar{x}$$
 (no nitrogen) = 0.399 gram,
 $mean \, \tilde{x}$ (no nitrogen) = $\frac{0.38 + 0.42}{2} = 0.400$ gram,
 $avg \, \bar{x}$ (nitrogen) = 0.565 gram,
 $mean \, \tilde{x}$ (nitrogen) = $\frac{0.49 + 0.52}{2} = 0.505$ gram.




Figure 1.4: Sample mean as a centroid of the with-nitrogen stem weight.







Figure 1.1: A dot plot of stem weight data.







Other Measures of Locations

Mignelide reini amidi (3)

-Trimmed Mean

e.g., in computing 10% trimmed mean, we cancel the highest 10% and the lowers 10% of our data TV % 10 -Benefit:

1) Having a mean close to median


2) Reduce the effect of very high and very low


value size is 10 for each sample. So for the without-nitrogen group the 10% trimmed mean is given by

$$\bar{x}_{\mathrm{tr}(10)} = \frac{0.32 + 0.37 + 0.47 + 0.43 + 0.36 + 0.42 + 0.38 + 0.43}{8} = 0.39750,$$

and for the 10% trimmed mean for the with-nitrogen group we have

$$\frac{14\bar{x}_{\text{tr}(10)}}{8} = \frac{0.43 + 0.47 + 0.49 + 0.52 + 0.75 + 0.79 + 0.62 + 0.46}{8} = 0.56625.$$





Accyvace itive 1) median
abyl 2) trimned w
abyl 3) aug (med Vo (mean) 1 stri vimme 6 mass simball redigh (ol) Will filtration can outliers 11 alliques

			** Trimmed Mean for No-Nitrogen		
	No-Nitrogen	Nitrogen			
	0.28	0.26			
	0.32	0.43	$x_{\text{tr}(10)} = 0.32 + 0.36 + 0.37 + 0.38 + 0.42 + 0.43 + 0.43 + 0.47$		
	0.36	0.46	8		
	0.37	0.47	= 0.3975		
	0.38	0.49	-0.5575		
	0.42	0.52			
	0.43	0.62			
	0.43	0.75	** Trimmed Mean for Nitrogen		
-	0.47	0.79			
	0.53	0.86	$x_{\text{tr}(10)} = 0.43 + 0.46 + 0.47 + 0.49 + 0.52 + 0.62 + 0.75 + 0.79$		
			8		
	, 1		- 0 56625		

The trimmed means are close to both the mean and median for the individual samples

• Trimmed mean is more insensitive to putliers than the sample mean but not as insensitive as the median.

• The sample median is a special case of the trimmed mean in which all of the sample data arteliminated apart from the middle one or two observations.

trimmed) 1 Grapha alla median 11 a Mean

Measures of Variability

- -Range
- Variance and Standard deviation

Range = $X_{\text{max}} - X_{\text{min}}$

The sample variance, denoted by
$$s^2$$
, is given by

$$s^2 = \sum_{i=1}^{n} \frac{(x_i - \bar{x})^2}{n-1}.$$
The sample standard deviation, denoted by s , is the positive square root of

The sample standard deviation, denoted by s, is the positive square root of s^2 , that is,

$$s = \sqrt{s^2}$$
.

imple 1.4: In an example discussed extensively in Chapter 10, an engineer is interested in testing the "bias" in a pH meter. Data are collected on the meter by measuring the pH of a neutral substance (pH = 7.0). A sample of size 10 is taken, with results

7.07 7.00 7.10 6.97 7.00 7.03 7.01 7.01 6.98 7.08.

The sample mean \bar{x} is given by

 $\bar{x} = \frac{7.07 + 7.00 + 7.10 + \dots + 7.08}{10} = \frac{7.0250}{10}$

The sample variance s^2 is given by

$$s^{2} = \frac{1}{9}[(7.07 - 7.025)^{2} + (7.00 - 7.025)^{2} + (7.10 - 7.025)^{2} + \cdots + (7.08 - 7.025)^{2}] = 0.001939.$$


As a result, the sample standard deviation is given by

$$s = \sqrt{0.001939} = 0.044.$$

So the sample standard deviation is 0.0440 with n-1=9 degrees of freedom.

17

Statistical Modeling, Scientific Inspection, and Graphical Diagnostics إرصاء وُقِعَى



At times the model postulated may take on a somewhat complicated form. Consider, for example, a textile manufacturer who designs an experiment where cloth specimen that contain various percentages of cotton are produced. Consider the data in Table

Table 1.3: Tensile Strength

deres cans allow to low seid


Descriptive.

1.3.

Statstics.

Cotton Percentage	Tensile Strength
15 %	7, 7, 9, 8, 10
20	19, 20, 21, 20, 22
25	21, 21, 17, 19, 20
30	8, 7, 8, 9, 10

terrile o chart penje bajans
terrile o chart penje bajans
serrength



tensile o miller (à citali ami au restali *
strength
(ainsies) tensile des chall sille des
19 aintes) strength

الغمن داورق 2) Stem-and-Leaf Plot

e.g., To illustrate the construction of a stem-and-leaf plot, consider the data of Table 1.4, which specifies the "life" of 40 similar car batteries recorded to the nearest tenth of a year.

	betsout to	Table 1.4: Car Battery Life					ife	75	,	
	[2.2]	4.1	3.5	4.5	3.2	3.7	3.0	2.6	L	
2.2	3.4	1.6	3.1	3.3	3.8	3.1	4.7	3.7	1	NO MARKET
it dear	3.3 3.3	4.3	3.4	3.6	2.9	3.3	3.9	3.1	DOE	25669
stork 1	3.3	3.1	3.7	4.4	3.2	4.1	1.9	3.4	7	2500/
	4.7	3.8	3.2	2.6	3.9	3.0	4.2	3.5		

data JIKin ax plia 1 (8 person stem 11 cus	Table 1.5: Stem-and-Leaf Plot of Battery Life					
1 100 1 Copy	Stem	Leaf	Frequency			
Stemila	1	69	2			
leaf 11 imps	2	25669 0011112223334445567778899	5 25			
1. でなる	4	11234577	8			
mean light		1	-> (1-4) (1) (1)			
20		1	* - (5-9)			